

THE CLASSIFICATION OF MATRICES

It is a fundamental result of Linear Algebra that, in any dimension n , there is a small number of simple ‘standard forms’ so that every $n \times n$ matrix is equivalent to one in standard form. ‘Equivalent’ is understood in the usual sense of the change of basis formula: if Λ is the standard form of A , there is an invertible matrix P so that $P^{-1}AP = \Lambda$. (Λ is uniquely defined given A -except for the order in which the eigenvalues appear- but many different P will work).

This notion of equivalence is natural- it means that, with respect to the basis of \mathbb{R}^n given by the columns of P , the action of A is very simple. In addition, reduction to standard form is an extremely useful procedure- for example, to solve systems of linear differential equations or difference equations.

The simplest situation occurs if A is diagonalizable with real eigenvalues- then the standard form is purely diagonal, with the eigenvalues themselves as diagonal entries. Allowing complex eigenvalues (that is, when \mathbb{C}^n has a basis consisting of eigenvectors of A) introduces the minor complication of 2×2 blocks on the diagonal, of the form $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$. When $n = 2$, there is only one other possibility (A has a double real eigenvalue and is not diagonalizable- example 1 below). When $n = 3$, there are 3 non-diagonalizable ‘standard forms’. The purpose of this handout is to describe the cases $n = 2, 3$ completely. The general case is dealt with in ‘advanced’ courses, but follows the same pattern. But keep in mind that, if you pick an $n \times n$ matrix ‘at random’, the probability of picking a non-diagonalizable one is ZERO. (It is like trying to hit $(0,0)$ by throwing darts at \mathbb{R}^2 - you’d have to be very unlucky.)

Example 1.

$$A = \begin{bmatrix} 3 & -1 \\ 1 & 1 \end{bmatrix}.$$

The characteristic polynomial is $p(\lambda) = (\lambda - 2)^2$, so 2 is the only eigenvalue. To compute its eigenspace, we solve the homogeneous system:

$$(A - 2I)v = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 0,$$

and obtain:

$$E(2) = \{t(1, 1); t \in \mathbb{R}\}.$$

So $E(2)$ is one-dimensional; since there are no other eigenvalues, A is not diagonalizable. To find a basis of \mathbb{R}^2 in which A acts in a simple way, we need another vector. We find one by picking an eigenvector (say, $v = (1, 1)$ itself) and solving the non-homogeneous system for w :

$$(A - 2I)w = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = v = \begin{bmatrix} 1 \\ 1 \end{bmatrix},$$

which is really the single equation $x - y = 1$. Any solution will do; take, say, $w = (1, 0)$. Then the action of A on the basis $\mathcal{B} = \{v, w\}$ is extremely simple:

$$Av = 2v; \quad Aw = 2w + v.$$

This means the matrix of A in this basis is:

$$[A]_{\mathcal{B}} = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix} := \Lambda.$$

This Λ is the ‘standard form’ of A . A matrix P satisfying $P^{-1}AP = \Lambda$ is given by the vectors of \mathcal{B} as columns: (*Watch the order!*)

$$P = [v|w] = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

Note that, although $(A - 2I)w \neq 0$ (since w is not an eigenvector), we do have:

$$(A - 2I)^2w = (A - 2I)(A - 2I)w = (A - 2I)v = 0,$$

since v is an eigenvector. w is said to be a ‘generalized eigenvector of order 2’.

The general fact behind the classification is that, even when there is no basis of \mathbb{R}^n consisting of eigenvectors of A , there is always a (computable!) basis consisting of *generalized* eigenvectors (that is v and λ satisfying $(A - \lambda I)^k v = 0$ for some $k \geq 0$ (k is at most n)). This very simple and general fact should be known to the average educated citizen, but usually gets buried under a mountain of impenetrable jargon in an ‘advanced’ course. (Behind the jargon are important concepts that make a proof for arbitrary n feasible.)

In summary, there are three ‘standard forms’ in the 2×2 case:

$$\begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}, \quad \begin{bmatrix} a & b \\ -b & a \end{bmatrix}, \quad \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}.$$

In the first one, A is diagonalizable with real eigenvalues (λ may be equal to μ , but only if A is just λ times the identity matrix, already in the standard basis). In the second, we take $b > 0$; A has a complex-conjugate pair of eigenvalues: $\lambda = a \pm ib$. The third case is non-diagonalizable: λ has algebraic multiplicity two (i.e., is a double root of the characteristic polynomial), but ‘geometric multiplicity’ 1 (dimension of the eigenspace). The corresponding matrices $P = [v|w]$ are easily computable: in the first case, v (resp. w) is an eigenvector for λ (resp. μ); in the second, v (resp. w) is the real part (resp. imaginary part) of an eigenvector for $a + ib$ (where $b > 0$). In the last case, v is an eigenvector for λ and w is any solution of the non-homogeneous system $(A - \lambda I)w = v$.

Moving on to $n = 3$, the diagonalizable (or complex-diagonalizable) cases are easy; their standard forms are:

$$\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \gamma \end{bmatrix}, \quad \begin{bmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & \gamma \end{bmatrix}.$$

In the first case, A is diagonalizable with real eigenvalues; in fact, two of λ, μ, γ may coincide, or even all three (if A is already of the form λ times the identity). In any case, the dimensions of the eigenspaces add up to 3. In the second, A has a real eigenvalue γ and a complex-conjugate pair of eigenvalues $a \pm ib$. The diagonalizing matrix P is given by $P = [v|w|u]$, where u is an eigenvector for γ and v (resp. w) is the real part (resp. imaginary part) of an eigenvector for $a + ib$ (where $b > 0$). Just as in two dimensions.

There are three non-diagonalizable cases (depending on how you count!) The first one has standard form:

$$\begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \mu \end{bmatrix}.$$

(We assume $\lambda \neq \mu$!) This is easily understood as a ‘non-diagonalizable 2×2 block’ for the eigenvalue λ , followed by a real eigenvalue μ . Here we have: $\dim(E(\lambda)) = \dim(E(\mu)) = 1$. The difference is that μ has algebraic multiplicity 1- it is a simple root of the characteristic polynomial-while λ is a double root (i.e., it has algebraic multiplicity 2). The ‘diagonalizing matrix’ $P = [v|w|u]$ consists of any eigenvectors v (resp. u) for λ (resp. μ), plus a ‘generalized eigenvector’ w for λ , satisfying $(A - \lambda I)w = v$. The action of A on the basis $\mathcal{B} = \{v, w, u\}$ is:

$$Av = \lambda v, \quad Aw = \lambda w + v, \quad Au = \mu u.$$

The two other cases are best explained using explicit examples.

Example 2.

$$A = \begin{bmatrix} -3 & 9 & 1 \\ -4 & 9 & 1 \\ 0 & 0 & 3 \end{bmatrix}.$$

Here the characteristic polynomial is $p(\lambda) = (\lambda - 3)^3$. So 3 is the only eigenvalue, with algebraic multiplicity 3. To compute its eigenspace, we solve the homogeneous system:

$$(A - 3I)v = \begin{bmatrix} -6 & 9 & 1 \\ -4 & 6 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 0.$$

The first two equations give:

$$-2x + 3y + (1/3)z = 0, \quad -2x + 3y + (1/2)z = 0,$$

so it follows that $z = 0$ and $2x = 3y$. Hence $E(3)$ is one-dimensional:

$$E(3) = \{t(3, 2, 0), t \in \mathbb{R}\}.$$

To get a basis for \mathbb{R}^3 we need two more vectors, which will be ‘generalized eigenvectors’ for $\lambda = 3$. We use the same method as when $n = 2$; pick a (non-zero) eigenvector (say, $v = (3, 2, 0)$ itself), and solve the non-homogeneous system:

$$(A - 3I)w = \begin{bmatrix} -6 & 9 & 1 \\ -4 & 6 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = v = \begin{bmatrix} 3 \\ 2 \\ 0 \end{bmatrix}.$$

This reduces to the equations:

$$-2x + 3y + (1/3)z = 1, \quad -2x + 3y + (1/2)z = 1,$$

so again it follows that $z = 0$, and x, y satisfy: $-2x + 3y = 1$ (*). Any solution of this will do; say, $x = y = 1$, leading to $w = (1, 1, 0)$.

Now, here you might think that one could get a third vector u simply by taking another (linearly independent) solution of (*)- say, $x = 0, y = 1/3$, but this won’t work. The reason is that any two pairs (x, y) solving (*) differ by a solution of $-2x + 3y = 0$, exactly the equation defining the eigenspace. So if we did this, $u - w$ would be a multiple of v , and the matrix $P = [v|w|u]$ would not be invertible. We have to try something else!

A natural thing to try is to simply repeat the procedure that generated w , that is, solve the non-homogeneous system:

$$(A - 3I)u = \begin{bmatrix} -6 & 9 & 1 \\ -4 & 6 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = w = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}.$$

This time we get:

$$-2x + 3y + (1/3)z = 1/3, \quad -2x + 3y + (1/2)z = 1/2,$$

and subtracting yields $(1/6)z = 1/6$, or $z = 1$. Then (x, y) can be *any* solution of $-2x + 3y = 0$, including $x = y = 0$, which gives $u = (0, 0, 1)$. The procedure worked: we have for the action of A on the basis $\mathcal{B} = \{v, w, u\}$:

$$Av = 3v, \quad Aw = 3w + v, \quad Au = 3u + w,$$

so the ‘standard form’ of A is:

$$\Lambda = [A]_{\mathcal{B}} = \begin{bmatrix} 3 & 1 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{bmatrix}.$$

The matrix $P = [v|w|u]$ brings A to standard form: $P^{-1}AP = \Lambda$. Note that w and u are ‘generalized eigenvectors’ of orders 2 and 3 (respectively):

$$(A - 3I)^3 u = (A - 3I)(A - 3I)(A - 3I)u = (A - 3I)(A - 3I)w = (A - 3I)v = 0.$$

There is one possibility left when $n = 3$.

Example 3.

$$A = \begin{bmatrix} -3 & 9 & 0 \\ -4 & 9 & 0 \\ -6 & 9 & 3 \end{bmatrix}.$$

Again, the characteristic polynomial is $p(\lambda) = (\lambda - 3)^3$, so $\lambda = 3$ is the only eigenvalue. As always, we begin by computing its eigenspace:

$$(A - 3I)v = \begin{bmatrix} -6 & 9 & 0 \\ -4 & 6 & 0 \\ -6 & 9 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 0.$$

This amounts to the single equation: $-2x + 3y = 0$ (with z unconstrained), so this time we have a two-dimensional eigenspace:

$$E(3) = \{s(3, 2, 0) + t(0, 0, 1); s, t \in \mathbb{R}\}.$$

We only need one more vector w to build a basis, and (given recent experience) to find it we solve the non-homogeneous system:

$$(A - 3I)w = \begin{bmatrix} -6 & 9 & 0 \\ -4 & 6 & 0 \\ -6 & 9 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = v,$$

where v is an eigenvector. Previously we've had essentially only one choice (up to constant multiples), but now we must face the question:

Which eigenvector do we pick?

The choice is not at all arbitrary, if we want our non-homogeneous system to be consistent. So pick a generic vector $v = s(3, 2, 0) + t(0, 0, 1)$ in $E(3)$, and try to solve:

$$\begin{bmatrix} -6 & 9 & 0 \\ -4 & 6 & 0 \\ -6 & 9 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = v = \begin{bmatrix} 3s \\ 2s \\ t \end{bmatrix}.$$

This leads to:

$$-2x + 3y = s, \quad -2x + 3y = s, \quad -2x + 3y = t/3,$$

so we need $s = t/3$ to get a consistent system. Any non-zero values will do; say $t = 3, s = 1$, which gives $v = (3, 2, 3)$, and the equation for w : $-2x + 3y = 1$ (z unconstrained). Any solution will do here- say, $x = y = 1, z = 0$, and then $w = (1, 1, 0)$.

Having picked $v \in E(3)$ and w satisfying $(A - 3I)w = v$, for u we may take any vector in $E(3)$ linearly independent with w - say, $u = (0, 0, 1)$. The action of A on the basis $\mathcal{B} = \{v, w, u\}$ is then:

$$Av = 3v, \quad Aw = 3w + v, \quad Au = 3u,$$

and we have the standard form for A :

$$\Lambda = [A]_{\mathcal{B}} = \begin{bmatrix} 3 & 1 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

(This looks like the first non-diagonalizable 3×3 case listed above before example 2, when $\lambda = \mu$, but it is best to count it separately). The matrix P taking A to standard form is (of course) $P = [v|w|u]$, and $P^{-1}AP = \Lambda$.

So, to complete the list, the two 3×3 non-diagonalizable standard forms just described are:

$$\begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}, \quad \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}.$$

The dimensions of the eigenspaces are: $\dim E(\lambda)=1$ in the first case, $\dim E(\lambda) = 2$ in the second. The algebraic multiplicity of λ is 3 in both cases.

EXERCISES. This topic is not difficult, but you won't learn it unless you get some practice. So here are **10** problems for you. In each case, a matrix A is given, and also its eigenvalues (listed with *algebraic* multiplicities). Your mission (which you would choose not to accept at your peril) is to find the standard form in each case, as well as the matrix that reduces A to its standard form.

$$1. \begin{bmatrix} 3 & 4 \\ -1 & 7 \end{bmatrix} \quad \text{eigenvalues: } 5, 5.$$

$$2. \begin{bmatrix} 4 & -2 \\ 1 & 1 \end{bmatrix} \quad \text{eigenvalues: } 2, 3.$$

$$3. \begin{bmatrix} -7 & 15 \\ -6 & 11 \end{bmatrix} \quad \text{eigenvalues: } 2 \pm 3i.$$

$$4. \begin{bmatrix} -4 & 9 & 0 \\ -4 & 8 & 0 \\ 0 & 0 & 5 \end{bmatrix} \quad \text{eigenvalues: } 2, 2, 5.$$

$$5. \begin{bmatrix} -4 & 9 & 0 \\ -6 & 11 & 0 \\ 0 & 0 & 2 \end{bmatrix} \quad \text{eigenvalues: } 2, 2, 5.$$

$$6. \begin{bmatrix} 1 & 12 & 0 \\ -1 & 8 & 0 \\ 1 & -3 & 4 \end{bmatrix} \quad \text{eigenvalues: } 4, 4, 5.$$

$$7. \begin{bmatrix} 1 & 12 & 0 \\ -1 & 8 & 0 \\ 0 & 0 & 5 \end{bmatrix} \quad \text{eigenvalues: } 4, 5, 5.$$

8. $\begin{bmatrix} -3 & 24 & 12 \\ -2 & 11 & 4 \\ 4 & -16 & 5 \end{bmatrix}$ eigenvalues: $3, 5 \pm 4i$.

9. $\begin{bmatrix} 7 & -9 & 0 \\ 1 & 1 & 0 \\ -1 & 4 & 4 \end{bmatrix}$ eigenvalues: $4, 4, 4$.

10. $\begin{bmatrix} 4 & 0 & 3 \\ 0 & 4 & 1 \\ 0 & 0 & 4 \end{bmatrix}$ eigenvalues: $4, 4, 4$.

Answers. Note that (when $n = 2$ or 3) to find the standard form it is enough to know the dimensions of the eigenspaces; this is false in dimensions 4 and higher. The matrices below are given by **rows**, in the following notation:

$$[[a, b][c, d]] = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

Recall that there is a unique correct answer for Λ , except for the order of the eigenvalues along the diagonal (which affects the order of the columns of P), but many possible correct answers for P , other than the one given. The first matrix given is Λ ; the second is P .

1. $[[5, 1][0, 5]][[2, -1][1, 0]]$
2. $[[2, 0][0, 3]][[1, 2][1, 1]]$
3. $[[2, 3][-3, 2]][[5, 0][3, 1]]$
4. $[[2, 1, 0][0, 2, 0][0, 0, 5]][[3, 1, 0][2, 1, 0][0, 0, 1]]$
5. $[[2, 0, 0][0, 2, 0][0, 0, 5]][[3, 0, 1][2, 0, 1][0, 1, 1]]$
6. $[[4, 1, 0][0, 4, 0][0, 0, 5]][[0, 4, 3][0, 1, 1][1, 0, 0]]$
7. $[[5, 0, 0][0, 5, 0][0, 0, 4]][[3, 0, 4][1, 0, 1][0, 1, 0]]$
8. $[[5, 4, 0][-4, 5, 0][0, 0, 3]][[-12, 13, 4][-3, 1, 1][1, 0, 0]]$
9. $[[4, 1, 0][0, 4, 1][0, 0, 4]][[0, 3, 4][0, 1, 1][1, 0, 0]]$
10. $[[4, 1, 0][0, 4, 1][0, 0, 4]][[3, 0, 1][1, 1, 1][0, 0, 1]]$