

TOPOLOGY PRELIM REVIEW 2021: LIST SIX

Topic: connected spaces.

1. A subset $E \subset \mathbb{R}$ is connected iff E is an interval.

2. (i) If each X_α is connected and $\bigcap_\alpha X_\alpha \neq \emptyset$, then $X = \bigcup_\alpha X_\alpha$ is connected.

(ii) If each pair of points $x, y \in X$ lie in some connected subset $E_{xy} \subset X$, then X is connected.

(iii) If $X = \bigcup_{n \geq 1} X_n$, each X_n is connected and $X_n \cap X_{n+1} \neq \emptyset \forall n \geq 1$, then X is connected.

3. If $E \subset X$ is connected and $E \subset A \subset \overline{E}$, then A is connected (in particular, \overline{E} is connected.)

4. $X = X_1 \times X_2$ is connected iff both X_i are.

Def. A chain of open sets connecting two points $a, b \in X$ is a sequence U_1, \dots, U_n of open sets of X such that $a \in U_1, b \in U_n$ and $U_i \cap U_j \neq \emptyset$ iff $|i - j| \leq 1$.

5. If X is connected and \mathcal{U} is any open cover of X , then any two points $a, b \in X$ can be connected by a chain of open sets consisting of elements of \mathcal{U} .

Hint: Let Z be the set of points of X that can be connected to a by a chain of open sets in \mathcal{U} . It is enough to show Z is closed (since it is clearly open and non-empty.)

6. If $E \subset \mathbb{R}^n$ is countable, its complement E^c is path connected.

7. A connected, locally path connected space X is path connected. In particular, connected open sets in \mathbb{R}^n are path connected.

8. The unit sphere in a normed vector space (of dimension at least 2) is path connected.

Def. X is *locally connected* if each $x \in X$ has a local basis of open connected sets.

9. X is locally connected iff each component of an open set is open.

10. If $f : X \rightarrow Y$ is a continuous map, and X is connected, then its graph $\Gamma_f \subset X \times Y$ is connected. Is the converse true?

11. (i) Any collection \mathcal{U} of pairwise disjoint open subsets of \mathbb{R}^n is countable. (In particular, any open subset of \mathbb{R}^n is the disjoint union of countably many connected components, or path components.)

(ii) If $I \subset \mathbb{R}$ is an interval and $f : I \rightarrow \mathbb{R}$ is a monotone function, the set of points of discontinuity of f is a countable subset of I .

(iii) Any open subset of \mathbb{R} can be expressed (in a unique way) as a countable union of disjoint open intervals.

12. A connected metric space with more than one point is uncountable. Hence any countable metric space is totally disconnected.

13. Let $f : X \rightarrow Y$ be a continuous, open, surjective map. If Y is connected and each fiber $f^{-1}(y)$ is connected, then X is connected. What about “path connected”?

Def. A space X is *totally disconnected* if its connected components are points.

14. (i) A subset $E \subset \mathbb{R}$ is totally disconnected iff E contains no interval.

(ii) $\mathbb{Q}, \mathbb{P} = \mathbb{R} \setminus \mathbb{Q}$ and the Cantor middle-thirds set $C \subset [0, 1]$ are totally disconnected.