

On the Geometric Convergence of Fractional Invariants in Non-Euclidean Topologies

Joe Random¹ Jane Smith² Robert Jones^{†2}

¹*Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA*

²*Department of Mathematics, Vanderbilt University, Nashville, TN 37240, USA*

Received: January 15, 2026; Revised: April 20, 2026; Accepted: May 10, 2026

Abstract. *This paper establishes an analytical framework for calculating the geometric convergence rates of fractional invariant measures on complete Riemannian manifolds. By extending classical localization techniques to localized fractional laplacians, we prove that the structural stability of topological entropy remains invariant under bounded metric perturbations. Our findings generalize several foundational theorems in non-Euclidean dynamics and provide new algorithmic lower bounds for multi-scale numerical integration schemes.*

1 Introduction

Let $(\mathcal{M}, g)$ be a complete, smooth Riemannian manifold with non-positive sectional curvature. The study of invariant measures under the action of fractional diffusion operators has seen significant development. In this paper, we address the open problem regarding structural stability limits when the metric g undergoes local variations.

The layout of this template tests your exact geometry choices (1.2 in margins, 0.75 in vertical buffers). Notice that subsequent page headers automatically display the parsed author string or short title without clipping into the printer's boundary box.

2 Core Theoretical Results

We begin by establishing the necessary analytical bound requirements to verify that your class file handles structural mathematical formulas alongside standard mathematical theorem counters cleanly.

2020 Mathematics Subject Classification. 54A05, 58J20, 37C40

Keywords: Fractional invariants, geometric convergence, Riemannian manifolds, topological entropy

E-mail addresses: JRandom@utk.edu (J. Random), jsmith@ornl.gov, rjones@gmail.com (R. Jones), and jsmith@vanderbilt.edu (J. Smith)

[†] Corresponding author

Definition 2.1. A fractional invariant measure μ is said to be geometrically regular if there exists a localized tracking constant $\Lambda > 0$ such that for all test functions $\phi \in C_c^\infty(\mathcal{M})$:

$$\int_{\mathcal{M}} (-\Delta)^\alpha \phi d\mu \leq \Lambda \|\phi\|_{L^2(\mathcal{M})}. \quad (1)$$

With our basic definitions established, we present the principal limiting convergence theorem below.

Theorem 2.2 (Main Convergence Property). *Let μ_n be a sequence of geometrically regular fractional invariant measures on $(\mathcal{M}, g)$. If the structural tracking capacity satisfies $\lim_{n \rightarrow \infty} \Lambda_n = \Lambda_\infty$, then μ_n converges weakly to a limiting measure μ_∞ satisfying the metric condition:*

$$|\mu_n(E) - \mu_\infty(E)| = \mathcal{O}\left(n^{-\frac{2\alpha}{d}}\right). \quad (2)$$

Proof. The proof proceeds via standard localization techniques. Let $\Omega \subset \mathcal{M}$ be a compact subset. By applying the fractional Sobolev embedding theorem directly to our kernel matrix, we deduce that the local trace operator is completely bounded. The remaining convergence constraints follow immediately from standard density arguments. $\square$

Lemma 2.3. *Under the assumptions of Theorem 2.2, any bounded perturbation of the background manifold metric g preserves weak compactness properties.*

Remark 2.4. Note that if $\alpha \rightarrow 1$, the convergence rate smoothly degenerates back into the classical local operator bounds established by classical diffusion models.

3 Numerical Verification and Benchmarking

To verify how multi-column floats or layout matrices interact with your document text block width (6.1 in), numerical test suites were compiled using local cluster instances. The resulting text flows cleanly around the mathematical frameworks.

Acknowledgments

J. Random was supported by NSF, grant #123456789. The authors express their gratitude to Dr. X. Fang and the University of Tennessee for the hospitality. Computing resources were graciously provided by the Advanced Scientific Computing Research infrastructure cluster.

References

- [1] T. Nguyen, *An Advanced Proof in Topology*, Univ. Tenn. Math. J. **5** (2026), no. 2, 12–45.
- [2] J. Smith, *Foundations of Non-Euclidean Geometry*, Academic Press, New York, 2024.

- [3] R. Jones and S. Wise, *Fractional Laplacian Perturbations on Riemannian Manifolds*, Journal of Functional Analysis, **288** (2025), no. 4, 102–129.
- [4] E. Özsoy, *Geophysical Fluid Dynamics I: An Introduction to Atmosphere - Ocean Dynamics: Homogeneous Fluids*, Springer Textbooks in Earth Sciences, Geography and Environment (2020).
- [5] J. Pedlosky, *Geophysical Fluid Dynamics*. Society for Industrial and Applied Mathematics (2005).
- [6] M. Sammartino and R. E. Caflisch, *Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a half-space. I. Existence for Euler and Prandtl equations*. Comm. Math. Phys. 192 (1998), no. 2, 433-461.
- [7] M. Sammartino and R. E. Caflisch, *Zero viscosity limit for analytic solutions of the Navier-Stokes equation on a half-space. II. Construction of the Navier-Stokes solution*. Comm. Math. Phys. 192 (1998), no. 2, 463-491.
- [8] H. Swann, *The convergence with vanishing viscosity of non-stationary Navier-Stokes flow to ideal flow in $\mathbb{R}^3$* , Trans. Amer. Math. Soc. 157 (1971), 373-397.
- [9] A. Tarfulea, *Improved a priori bounds for thermal fluid equations*. Trans. Amer. Math. Soc. 371 (2019), no. 4, 2719-2737.
- [10] R. Temam, *On the Euler equations of incompressible perfect fluids*, J. Func. Anal. 20 (1975), 32-43.
- [11] R. Temam, *Navier-Stokes Equations*, North-Holland, 1983.
- [12] F. Wang, *The three-dimensional inviscid limit problem with data analytic near the boundary*. SIAM J. Math. Anal. 52 (2020), no. 4, 3520-3545.