

TIME-DEPENDENCY IN HYPERBOLIC ANDERSON MODELS: STRATONOVICH REGIME

BY XIA CHEN^{1,a},

¹*Department of Mathematics, University of Tennessee, Knoxville, Tennessee, USA*, ^axchen3@tennessee.edu

In this paper, the hyperbolic Anderson equation generated by a time-dependent Gaussian noise is under investigation in two fronts: The solvability and large- t asymptotics. The investigation leads to a necessary and sufficient condition for existence and a large- t limit form for the expectation of the solution. Three major developments are made for achieving these goals: A universal bound for Stratonovich moment that guarantees the Stratonovich integrability and $\mathcal{L}^2$ -convergence of the Stratonovich chaos expansion under the best possible condition, a representation of the expected Stratonovich moments in terms of a time-randomized Brownian intersection local time, and a large deviation principle for the time-randomized Brownian intersection local time.

CONTENTS

1	Introduction	1
2	Stratonovich expansion and approximations	7
2.1	Green's function	7
2.2	Smoothed version of $\dot{W}(t, x)$	7
2.3	Stratonovich integral	9
3	Stratonovich moment bound	14
4	Stratonovich integrability of $g_n(\cdot, t, x)$	22
5	Proof of Theorem 1.1	33
6	A Stratonovich moment representation	37
7	The time-randomized intersection local time	42
8	Proof of Theorem 1.2	55
8.1	Upper bound for (8.1)	56
8.2	Lower bound for (8.1)	58
	Appendix	60
	Funding	63
	References	63

1. Introduction. The model studied in this paper is named after the physicist Philip Warren Anderson ([1]) who studies the random Schrödinger operator $\Delta + \dot{W}$ in his investigation of magnetic impurities embedded in metals. When the random Schrödinger operator appears in the heat equation (see (1.6) below), the resulting model is called parabolic Anderson model ([18]). We refer the interested readers to [22] and the references therein for general information on parabolic Anderson models, and to [26] for discrete versions of parabolic Anderson models.

Due to its close links to other physical models such as KPZ equation ([28]), especially in the wake of the breakthrough of [25], the study of this equation has been rapidly developed.

MSC2020 subject classifications: Primary 60F10, 60H15, 60H40, 60J65; Secondary 81U10.

Keywords and phrases: hyperbolic and parabolic Anderson equations, Stratonovich integrability, multiple Stratonovich integrals, multiple Stratonovich integrals, Brownian motion, time-randomized intersection local time, Wick's formula, moment asymptotics, intermittency.

The development is also credited to the Feynman-Kac formula which links the solution to some functionals of Brownian motions

When the Schrödinger operator appears the wave equation (see (1.1) below), the model is known as hyperbolic Anderson model. Due to lack of analytic tools such as Feynman-Kac formula, much less have been known on hyperbolic models. Until very recent, it was widely believed that as $\partial u/\partial t$ being replaced by $\partial^2 u/\partial t^2$, the hyperbolic equation has a wilder behavior than its parabolic counterpart. The recent progress shows otherwise: The hyperbolic equations are solvable under the conditions ([4], [11]) generally weaker than those posted for the existence of parabolic systems in the same regime.

It is also noticeable that most of the recent development cited above focus on the setting of time-independent Gaussian noise. The reason behind is that the time-dependency of the Gaussian field posts serious challenges and limitations to the currently available tools and ideas in dealing with hyperbolic Anderson models. We shall see a fundamental difference on the representation of the Stratonovich moment between the settings of the time-independent noise (Theorem 3.3, [14]) and the time-dependent noise (Theorem 6.1 below).

In this paper we consider the hyperbolic Anderson equation

$$(1.1) \quad \begin{cases} \frac{\partial^2 u}{\partial t^2}(t, x) = \Delta u(t, x) + \dot{W}(t, x)u(t, x), & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d \\ u(0, x) = 1 \text{ and } \frac{\partial u}{\partial t}(0, x) = 0, & x \in \mathbb{R}^d \end{cases}$$

run by a mean zero and possibly generalized Gaussian noise $\dot{W}(t, x)$ fractional in time, defined by its covariance function

$$(1.2) \quad \text{Cov}(\dot{W}(s, x), \dot{W}(t, y)) = |s - t|^{-\alpha_0} \gamma(x - y), \quad s, t \in \mathbb{R}_+, \quad x, y \in \mathbb{R}^d,$$

where $0 < \alpha_0 < 1$. As a covariance function, the non-negative definiteness of $\gamma(\cdot)$ implies that it admits a spectral measure $\mu(d\xi)$ on $\mathbb{R}^d$ uniquely defined by the relation

$$(1.3) \quad \gamma(x) = \int_{\mathbb{R}^d} e^{i\xi \cdot x} \mu(d\xi), \quad x \in \mathbb{R}^d.$$

Throughout this work, we assume that $\gamma(\cdot) \geq 0$ and $d = 1, 2, 3$ and interpret the system in the sense Stratonovich. Roughly speaking, it means that the equation (1.1) is the result of the approximation by classical wave equations run by the smoothed Gaussian noise $\dot{W}_{\epsilon, \delta}(t, x)$. We shall provide the details of the construction of the solution in Section 2.

Our first concern is to find the best condition for the existence of solution. The conditions for solvability are often formulated in term of the integrability of the spectral measure $\mu(d\xi)$. In a different set-up known as Skorohod regime, Chen, Deya, Song and Tindel ([12]) posted the optimal condition

$$(1.4) \quad \int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^{\frac{3-\alpha_0}{2}} \mu(d\xi) < \infty$$

for the existence/uniqueness of the system (1.1).

In their follow-up paper [11] for Stratonovich regime, (1.1) is solved under an apparently less optimal assumption in the dimensions $d = 1, 2$.

The following is the first main theorem in this paper.

THEOREM 1.1. *Let $d = 1, 2, 3$.*

(1) *Under the condition*

$$(1.5) \quad \int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^{\frac{2-\alpha_0}{2}} \mu(d\xi) < \infty$$

the equation (1.1) has a solution in the sense of Definition 2.1 given in Section 2.

(2) *If the equation (1.1) has a square integrable solution $u(t, x)$ that admits the Stratonovich expansion (see (2.13)) for some $t > 0$, then condition (1.5) must be satisfied.*

For the purpose of comparison, we introduce the parabolic Anderson model

$$(1.6) \quad \begin{cases} \frac{\partial u}{\partial t}(t, x) = \Delta u(t, x) + \dot{W}(t, x)u(t, x), & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d \\ u(0, x) = 1 & x \in \mathbb{R}^d. \end{cases}$$

It is well known that the condition (see, e.g. Theorem 4.6, [33]) for (1.6) to be solvable (in a sense same as Theorem 1.1) is

$$(1.7) \quad \int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^{1-\alpha_0} \mu(d\xi) < \infty.$$

In other words, the hyperbolic Anderson model is solvable under a condition genuinely weaker than the one set for its parabolic counterpart when it comes to the Gaussian field that is fractional in time.

The next main theorem is about the increasing rate of $\mathbb{E}u(t, x)$ as $t \rightarrow \infty$. To this end, we assume the homogeneity for the covariance structure:

$$(1.8) \quad \gamma(cx) = c^{-\alpha} \gamma(x), \quad x \in \mathbb{R}^d \quad c > 0$$

for some $\alpha > 0$. Taking $f(\lambda) = (1 + \lambda^2)^{-\frac{2-\alpha_0}{2}}$ and $v(d\xi) = \mu(d\xi)$ in [12, Lemma 3.10] yields

$$\int_{\mathbb{R}^d} \left(\frac{1}{1 + |\xi|^2} \right)^{\frac{2-\alpha_0}{2}} \mu(d\xi) = \alpha \mu\{\xi \in \mathbb{R}^d; |\xi| \leq 1\} \int_0^\infty \left(\frac{1}{1 + \rho^2} \right)^{\frac{2-\alpha_0}{2}} \frac{d\rho}{\rho^{1-\alpha}}$$

as far as either side is finite. This shows that under the homogeneity (1.8) on the noise covariance condition, the condition (1.5) becomes “ $\alpha_0 + \alpha < 2$ ”. In addition (Remark 1.4, [12]), the fact that $\gamma(\cdot)$ is non-negative and non-negative definite (for being qualified as covariance function) requires that $\alpha \leq d$. Further, the only setting where “ $\alpha = d$ ” is allowed under $\alpha < 2$ is when $\alpha = d = 1$, or when $\gamma(\cdot)$ is a constant multiple of the Dirac function.

THEOREM 1.2. *Under the homogeneity condition (1.8) with*

$$(1.9) \quad \alpha_0 + \alpha < 2$$

with $0 < \alpha_0 < 1$ and $0 < \alpha < d$ or with $0 < \alpha_0 < 1$ and $\alpha = d = 1$,

$$(1.10) \quad \begin{aligned} & (3 - \alpha) \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4-\alpha-\alpha_0}} \left(\frac{2\mathcal{E}_*}{2 - \alpha} \right)^{\frac{2-\alpha}{2}} \right)^{\frac{1}{3-\alpha}} \\ & \leq \liminf_{t \rightarrow \infty} t^{-\frac{4-\alpha-\alpha_0}{3-\alpha}} \log \mathbb{E}u(t, x) \leq \limsup_{t \rightarrow \infty} t^{-\frac{4-\alpha-\alpha_0}{3-\alpha}} \log \mathbb{E}u(t, x) \\ & \leq (3 - \alpha) \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4-\alpha-\alpha_0}} \left(\frac{2\mathcal{E}}{2 - \alpha} \right)^{\frac{2-\alpha}{2}} \right)^{\frac{1}{3-\alpha}} \end{aligned}$$

where

$$(1.11) \quad \mathcal{E} = \sup_{g \in \mathcal{A}_d} \left\{ \int_0^1 \int_0^1 \int_{\mathbb{R}^d \times \mathbb{R}^d} \frac{\gamma(x-y)}{|s-r|^{\alpha_0}} g^2(s,x) g^2(r,y) dx dy ds dr \right. \\ \left. - \int_0^1 \int_{\mathbb{R}^d} |\nabla_x g(s,x)|^2 dx ds \right\}$$

$$(1.12) \quad \mathcal{E}_* = \sup_{g \in \mathcal{A}_d} \left\{ \int_0^1 \int_0^1 \int_{\mathbb{R}^d \times \mathbb{R}^d} \gamma_*^0(s-r) \gamma(x-y) g^2(s,x) g^2(r,y) dx dy ds dr \right. \\ \left. - \frac{1}{2} \int_0^1 \int_{\mathbb{R}^d} |\nabla_x g(s,x)|^2 dx ds \right\}$$

and where

$$\mathcal{A}_d = \left\{ g(s,x); \quad g(s,\cdot) \in W^{1,2}(\mathbb{R}^d) \quad \text{and} \quad \int_{\mathbb{R}^d} |g(s,x)|^2 dx = 1 \text{ for every } 0 \leq s \leq 1 \right\},$$

$$(1.13) \quad \gamma_*^0(u) = \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+} \exp \left\{ - \left(\lambda + \frac{\lambda^2}{2(4-\alpha-2\alpha_0)} \right) |u| \right\} \frac{d\lambda}{\lambda^{1-\alpha_0}}.$$

First, $\mathcal{E} < \infty$ under $\alpha < 2$ (Lemma 5.2, [8]). Second, comparing (1.13) with (1.21) below, it is not hard to see that $\mathcal{E}_* < \mathcal{E}$. The gap in (1.11) reflects a new challenge (see Remark 8.1 below) on the large- t asymptotics for the complex-valued intersection local time

$$(1.14) \quad \int_0^t \int_0^t \left(((s-r) + i(\beta(s) - \beta(r))) \operatorname{sgn}(s-r) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr$$

driven by the independent Brownian motions $\beta(t)$ (1-dimensional) and $B(t)$ (d -dimensional). The relevance of (1.14) to our model in (1.1) is given in Theorem 6.1 below. To close the gap we call for establishment of the precise large deviation principle for the logarithmic-exponential moment

$$\log \mathbb{E}_0 \exp \left\{ \frac{1}{t^{1-\alpha_0}} \int_0^t \int_0^t \left(((s-r) + i(\beta(s) - \beta(r))) \operatorname{sgn}(s-r) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\}$$

as $t \rightarrow \infty$. Noticed from (8.7) below that the exponential moment $\mathbb{E}_0 \exp\{\dots\}$ remains positive despite that the Hamiltonian in (1.14) takes complex values. Therefore, the proposed large deviation principle makes mathematical sense. On the other hand, some new idea is needed to handle the the large deviations for the complex-valued systems. Beyond technicality, we predict that the limit in (1.10) exists and is given by the upper bound in (1.10). This prediction is based on the belief (see the discussion in Section 7) that in the game of large deviations, the complex Hamiltonian in (1.14) behaves like the real intersection local time

$$\int_0^t \int_0^t |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr$$

A result close to (1.11) found in literature is obtained by Balan and Conus (Theorem 2.1, [3]) where the system (1.1) is set up in Skorohod regime and the bounds

$$\limsup_{t \rightarrow \infty} t^{-\frac{4-\alpha-\alpha_0}{3-\alpha}} \log \mathbb{E} |u(t,x)|^p \leq \frac{4-p}{3-p} C_2 < \infty \quad (p \geq 2)$$

(with $C_2 > 0$ independent of p) and

$$\liminf_{t \rightarrow \infty} t^{-\frac{4-\alpha-\alpha_0}{3-\alpha}} \log \mathbb{E} u^2(t, x) > 0$$

are obtained.

In the case of the parabolic Anderson model (1.6), the moment asymptotics (Theorem 6.1, [13]) follow the pattern

$$(1.15) \quad \lim_{t \rightarrow \infty} t^{-\frac{4-\alpha-2\alpha_0}{2-\alpha}} \log \mathbb{E} u^p(t, x) \quad p = 1, 2, \dots$$

under the scaling property (1.8)) with

$$(1.16) \quad 2\alpha_0 + \alpha < 2.$$

Comparing (1.10) and (1.9) with (1.15) and (1.16), respectively, one can see the contribution from the time component of the Gaussian field are different between hyperbolic and parabolic settings. See Remark 6.2 below for an explanation from a new perspective.

In the recent work ([14]) on the hyperbolic Anderson model with time-independent noise $\dot{W}(x)$, the solvability is established under the Dalang's condition

$$(1.17) \quad \int_{\mathbb{R}^d} \frac{1}{1 + |\xi|^2} \mu(d\xi) < \infty$$

and the long-term moment asymptotics are established in the form of

$$(1.18) \quad \lim_{t \rightarrow \infty} t^{-\frac{4-\alpha}{3-\alpha}} \log \mathbb{E} u^p(t, x) \quad p = 1, 2, \dots$$

under the condition $\alpha < 2$.

On the side of idea development, the current work is partially motivated by [14] where the Stratonovich moment is represented in terms of the intersection local times

$$\int_0^t \int_0^t \gamma(B(s) - B(r)) ds dr \quad \text{and} \quad \int_0^t \int_0^t \gamma(B(s) - \tilde{B}(r)) ds dr$$

driven by the independent d -dimensional Brownian motions $B(t)$ and $\tilde{B}(t)$ (see Corollary 3.3, [14] for details). This connection is established on the simple fact ((3.6), [14]) that

$$(1.19) \quad \int_0^\infty e^{-\lambda t} G(t, x) dt = \frac{1}{2} \int_0^\infty \exp\left\{-\frac{\lambda^2}{2} t\right\} p(t, x) dt \quad (\lambda > 0)$$

where $p(t, x)$ is the Brownian semi-group

$$(1.20) \quad p(t, x) = (2\pi t)^{-d/2} \exp\left\{-\frac{|x|^2}{2t}\right\} \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$$

and where $G(t, x)$ is the fundamental solution (see (2.2) below) of the wave equation. Indeed, one of the crucial observations (Theorem 6.1) made in the current work is a link between the Stratonovich moment and the time-randomized intersection local time given (1.14). Accordingly, a large deviation principle (Theorem 7.1) for the time-randomized intersection local time is established.

We now mention some recent works on this subject for the comparison on methodology. Compared to our treatment, the approaches used in [2] and [11] are more analytic: In [2], Malliavin calculus and Hu-Meyer formula are applied to approximate the Stratonovich moment; while in [11], Strichartz estimation is used to bound the integrals in milled equation. The fact that the conditions given in [2] and [11] for existence are significantly more restrictive indicates that the hyperbolic equations do not correspond to analytic control well

enough due to their oscillation behaviors. In ([12]), the existence is achieved under the optimal condition for the hyperbolic Anderson models in the Skorohod regime. A similar idea used in [12] is time-exponentiation to the moment of chaos expansion. It should be pointed out that our representation of the moments of chaos in terms of the Brownian intersection local time can not be applied to the Skorohod moment at the current level of development, due to the reason same as the one given in Remark 6.3 below.

Another important task carried out in this paper is the legalization of Stratonovich expansion (2.13) under the best condition (1.5). To this end, a universal bound (Theorem 3.1) is established that is responsible for the Stratonovich integrability and $\mathcal{L}^2$ -convergence of the Stratonovich expansion. It should be pointed out that Stratonovich moment demands a level of technology higher than its Skorohod counter part.

This paper also brings a different idea on the treatment of the time-covariance $|\cdot|^{-\alpha_0}$ (introduced in (1.2)). There have been two ways in literature in handling $|\cdot|^{-\alpha_0}$. In the work [3] and some of its follow-up papers, the Hardy–Littlewood–Sobolev inequality is used (Lemma B.3, [3]) in the Skorohod regime to separate the time component. This strategy appears to be powerful in the parabolic setting. In the hyperbolic setting, it does not function as good as for the parabolic equations, as it is less adoptive to the oscillation behavior of wave operator. Another existing practice (e.g., [12]) is to perform the Fourier transform

$$|u|^{-\alpha_0} = C \int_{\mathbb{R}} e^{i\lambda u} \frac{d\lambda}{|\lambda|^{1-\alpha_0}} \quad u \in \mathbb{R}$$

to $|\cdot|^{-\alpha_0}$. The use of Fourier transformation has been popular and effective in parabolic setting (and hyperbolic/Skorohod setting as well). For the hyperbolic equation in the Stratonovich regime, it creates an annoying and un-controllable singularity. Instead, it is proposed in this work to use the Laplace transform

$$(1.21) \quad |u|^{-\alpha_0} = \Gamma(\alpha_0)^{-1} \int_0^\infty e^{-\lambda|u|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \quad u \in \mathbb{R}.$$

It should be pointed out that the time-dependency in the hyperbolic Anderson model is far from fully understood. More specifically, a deeper connection between the Stratonovich expansion and the Brownian intersection local times is very likely. The progress should lead to a fully understanding of the intermittency of the system. Despite of its incompleteness, on the other hand, Theorem 1.2 strongly indicates the most possible pattern of intermittency

$$(1.22) \quad \lim_{t \rightarrow \infty} t^{-\frac{4-\alpha-\alpha_0}{3-\alpha}} \log \mathbb{E}|u(t, x)|^p \\ = (3-\alpha)p^{\frac{4-\alpha}{3-\alpha}} \left(\frac{(4-\alpha-2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}}}{2^3(4-\alpha-\alpha_0)^{4-\alpha-\alpha_0}} \left(\frac{2\mathcal{E}}{2-\alpha} \right)^{\frac{2-\alpha}{2}} \right)^{\frac{1}{3-\alpha}}$$

for $p = 1, 2, \dots$. Here we post it as conjecture and leave it for future study.

Here is the organization of the paper. In next section (Section 2), we set a way to approximate the possibly generalized Gaussian field $\dot{W}(t, x)$, introduce the multiple Stratonovich integral and formally express the solution as Stratonovich expansion. In Section 3, we establish a universal bound for the moments in the Stratonovich expansion. Using this bound we install the Stratonovich integrability for the functions $g_n(\cdot, t, x)$ in Section 4 and establish the convergence of the Stratonovich expansion (and therefore prove Theorem 1.1) in Section 5. In Section 6, we establish a link between $\mathbb{E}S_{2n}(g_{2n}(\cdot, t, x))$ and the n -th moment of a time-randomized Brownian intersection local time under time exponentiation. In Section 7, we prove a large deviation principle for the time-randomized Brownian intersection local time. Using the results from Section 6 and 7, we prove Theorem 1.2 in Section 8. In the appendix, we conduct some elementary calculation for the bounds established in Section 3.

2. Stratonovich expansion and approximations. As usual by the Duhamel principle the mathematical definition of the hyperbolic Anderson equation (1.1) will be the following mild form

$$(2.1) \quad u(t, x) = 1 + \int_0^t \int_{\mathbb{R}^d} G(t-s, x-y) u(s, y) W(ds, dy),$$

where

(i) $G(t, x)$ is the fundamental solution defined by the deterministic wave equation

$$(2.2) \quad \begin{cases} \frac{\partial^2 G}{\partial t^2}(t, x) = \Delta G(t, x) \\ G(0, x) = 0 \quad \text{and} \quad \frac{\partial G}{\partial t}(0, x) = \delta_0(x), \quad x \in \mathbb{R}^d. \end{cases}$$

(ii) the stochastic integral on the right hand side of (2.1) is interpreted in the sense of Stratonovich (see discussion below for details).

2.1. Green's function. The fundamental solution $G(t, x)$ associated with (2.2) plays a key role in determining the behavior of the system (2.1). Let us recall some basic facts. Taking the Fourier transform in (2.2) we get the expression for the fundamental solution

$$(2.3) \quad \mathcal{F}(G(t, \cdot))(\xi) = \int_{\mathbb{R}^d} G(t, x) e^{i\xi \cdot x} dx = \frac{\sin(|\xi|t)}{|\xi|}, \quad (t, \xi) \in \mathbb{R}_+ \times \mathbb{R}^d$$

in its Fourier transform form. In particular,

$$(2.4) \quad \int_{\mathbb{R}^d} G(t, x) dx = \mathcal{F}(G(t, \cdot))(0) = t$$

In the dimensions $d = 1, 2, 3$, the fundamental solution $G(t, x)$ itself can be expressed explicitly as

$$(2.5) \quad G(t, x) = \begin{cases} \frac{1}{2} 1_{\{|x| \leq t\}} & d = 1 \\ \frac{1}{2\pi} \frac{1_{\{|x| \leq t\}}}{\sqrt{t^2 - |x|^2}} & d = 2 \\ \frac{1}{4\pi t} \sigma_t(dx) & d = 3, \end{cases}$$

where $\sigma_t(dx)$ is the surface measure on the sphere $\{x \in \mathbb{R}^3; |x| = t\}$. We limit our attention to $d = 1, 2, 3$ in this work because the treatment developed here requires $G(t, x) \geq 0$ (when $G(t, x)$ exists only in the form of distribution, it means that $(G(t, \cdot) * h)(x) \geq 0$ for sufficiently smooth probability density $h(x)$ on $\mathbb{R}^d$). A scaling property we frequently use (especially in the proof of Theorem 1.2) is

$$(2.6) \quad G(t, x) = t^{-(d-1)} G(1, t^{-1}x), \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^d.$$

2.2. Smoothed version of $\dot{W}(t, x)$. Generally speaking, the smoothed version of the generalized Gaussian field $\dot{W}(t, x)$ can be any family $\{\dot{W}_{\epsilon, \delta}(t, x); \epsilon, \delta > 0\}$ of mean-zero Gaussian fields on $\mathbb{R}_+ \times \mathbb{R}^d$, living in the same probability space $(\Omega, \mathcal{A}, \mathbb{P})$ as $W(t, x)$ does,

such that for each $\epsilon, \delta > 0$, $\dot{W}_{\epsilon, \delta}(t, x)$ is defined point-wise even with path-continuity (if needed), and of the form of covariance function

$$(2.7) \quad \text{Cov} \left(\dot{W}_{\epsilon, \delta}(s, x), \dot{W}_{\epsilon, \delta}(t, y) \right) = \gamma_{\delta}^0(s-t) \gamma_{2\epsilon}(x-y) \quad (s, x), (t, y) \in \mathbb{R}_+ \times \mathbb{R}^d$$

that satisfies

$$(2.8) \quad \begin{aligned} & \lim_{\epsilon, \delta \rightarrow 0^+} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} \gamma_{\delta}^0(s-t) \gamma_{\epsilon}(x-y) f(s, x) g(t, y) ds dt dx dy \\ &= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma(x-y) f(s, x) g(t, y) ds dt dx dy \quad f, g \in \mathcal{S}(\mathbb{R}_+ \times \mathbb{R}^d) \end{aligned}$$

where $\mathcal{S}(\mathbb{R}_+ \times \mathbb{R}^d)$ is the Schwartz space of all functions on $\mathbb{R}_+ \times \mathbb{R}^d$ that are infinitely differentiable and rapidly decay to zero at infinity. We shall see the choice we made for $\gamma_{\delta}^0(\cdot)$ and $\gamma_{2\epsilon}(\cdot)$ in (2.9) below:

A popular construction of $\dot{W}_{\epsilon, \delta}$ in literature is by convolution:

$$\dot{W}_{\epsilon, \delta}(t, x) = \int_{\mathbb{R}^{d+1}} p_0(\delta, t-u) p(\epsilon, x-y) \dot{W}(u, y) du dy \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$$

where $p(t, x)$ is the Brownian semi-group defined on $\mathbb{R}^d$ that is given in (1.20) and $p_0(t, x)$ is the Brownian semi-group on $\mathbb{R}$ and where on the right hand side, the possibly generalized Gaussian noise $\dot{W}(\cdot, \cdot)$ is interpreted as a linear functional acting on the smooth function $p_0(\delta, t-\cdot) p(\epsilon, x-\cdot)$.

To promote the use of the Laplace transform proposed in (1.21), we shall construct $\dot{W}_{\epsilon, \delta}(t, x)$ in a slightly different way such that

$$(2.9) \quad \text{Cov} \left(\dot{W}_{\epsilon, \delta}(s, x), \dot{W}_{\epsilon, \delta}(t, y) \right) = \left(\Gamma(\alpha_0)^{-1} \int_0^{\delta^{-1}} e^{-\lambda|s-t|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \right) \gamma_{2\epsilon}(x-y)$$

for $(s, x), (t, y) \in \mathbb{R}_+ \times \mathbb{R}^d$, where

$$\gamma_{\epsilon}(x) = \int_{\mathbb{R}^d} p(\epsilon, x-y) \gamma(y) dy \quad x \in \mathbb{R}^d.$$

If it exists (i.e., if they can live in the same probability space with $W(t, x)$), the family $\{\dot{W}_{\epsilon, \delta}(\cdot, \cdot); \epsilon, \delta > 0\}$ meets all requirements as the smoothed version of $\dot{W}(\cdot, \cdot)$: First, by the expression

$$(2.10) \quad \Gamma(\alpha_0)^{-1} \int_0^{\delta^{-1}} e^{-\lambda|s-t|} \frac{d\lambda}{\lambda^{1-\alpha_0}} = \Gamma(\alpha_0)^{-1} \mathbb{E}^{\kappa} \int_0^{\delta^{-1}} e^{i\lambda\kappa(s-t)} \frac{d\lambda}{\lambda^{1-\alpha_0}}$$

the function of $s-t$ on the left is non-negative definite and therefore qualified to be used as covariance function, where κ is a standard 1-dimensional Cauchy random variable.

Second, (2.8) holds in light of (1.21).

Third, by the relation

$$\begin{aligned} & \mathbb{E} \left(\dot{W}_{\epsilon, \delta}(s, x) - \dot{W}_{\epsilon, \delta}(t, y) \right)^2 \\ &= 2\Gamma(\alpha_0)^{-1} \left\{ \left(\int_0^{\delta^{-1}} \frac{d\lambda}{\lambda^{1-\alpha_0}} \right) \gamma_{\epsilon}(0) - \left(\int_0^{\delta^{-1}} e^{-\lambda|s-t|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \right) \gamma_{\epsilon}(x-y) \right\} \\ &\leq C_{\epsilon, \delta} \left\{ |s-t| + |x-y| \right\} \end{aligned}$$

and therefore by normality

$$\mathbb{E} \left(\dot{W}_{\epsilon, \delta}(s, x) - \dot{W}_{\epsilon, \delta}(t, y) \right)^{2n} \leq \frac{(2n)!}{2^n n!} C_{\epsilon, \delta}^n \left\{ |s - t| + |x - y| \right\}^n$$

for any integer $n \geq 1$. A standard use of the Kolmogorov continuity theorem (Theorem D.7, p.313, [5]), the Gaussian field $\dot{W}_{\epsilon, \delta}(t, x)$ has a continuous modification on $\mathbb{R}_+ \times \mathbb{R}^d$.

As for the family $\{\dot{W}_{\epsilon, \delta}; \epsilon, \delta > 0\}$ live in the same probability space, we start with the following simple observation: Given $0 < T_1 < \dots < T_n < \dots$, let $\Delta_k \dot{W}(t, x)$ ($k = 1, 2, \dots$) be independent mean-zero (possibly generalized) Gaussian fields on $\mathbb{R}_+ \times \mathbb{R}^d$ with the covariance functions

$$\text{Cov} \left(\Delta_k \dot{W}(s, x), \Delta_k \dot{W}(t, y) \right) = \left(\Gamma(\alpha_0)^{-1} \int_{T_{k-1}}^{T_k} e^{-\lambda|s-t|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \right) \gamma(x - y)$$

for $(s, x), (t, y) \in \mathbb{R}_+ \times \mathbb{R}^d$, where we follow the convention $T_0 = 0$. Set

$$\dot{W}_{T_n}(t, x) = \sum_{k=1}^n \Delta_k \dot{W}(t, x) \quad n = 1, 2, \dots$$

It is straightforward to see that

$$\text{Cov} \left(\dot{W}_{T_n}(s, x), \dot{W}_{T_n}(t, y) \right) = \left(\Gamma(\alpha_0)^{-1} \int_0^{T_n} e^{-\lambda|s-t|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \right) \gamma(x - y).$$

Without changing its distribution, one can re-define the Gaussian field $W(t, x)$ as

$$\dot{W}(t, x) = \sum_{k=1}^{\infty} \Delta_k \dot{W}(t, x)$$

for any monotonic sequence $\{T_n\}$ satisfying $T_n \rightarrow \infty$ ($n \rightarrow \infty$).

By Kolmogorov consistence extension, we can extend $\{\dot{W}_{T_n}(\cdot, \cdot); n = 1, 2, \dots\}$ to the bigger family $\{\dot{W}_T(\cdot, \cdot); T > 0\}$. Then we adopt the new notation that replaces $\dot{W}_{\delta^{-1}}(\cdot, \cdot)$ by $\dot{W}_{\delta}(\cdot, \cdot)$. Finally we define

$$\dot{W}_{\epsilon, \delta}(t, x) = \int_{\mathbb{R}^d} p(\epsilon, x - y) \dot{W}_{\delta}(t, y) dy$$

which satisfies (2.9). More generally

$$(2.11) \quad \text{Cov} \left(\dot{W}_{\epsilon, \delta}(s, x), \dot{W}_{\tilde{\epsilon}, \tilde{\delta}}(t, y) \right) = \gamma_{\delta \vee \tilde{\delta}}^0(s - t) \gamma_{\epsilon + \tilde{\epsilon}}(x - y)$$

for any $\epsilon, \tilde{\epsilon}, \delta, \tilde{\delta} > 0$.

2.3. Stratonovich integral. Given a random field $\{\Psi(t, x)\}_{(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d}$ such that

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} \Psi(t, x) \dot{W}_{\epsilon, \delta}(t, x) dt dx \in \mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P}) \quad \forall \epsilon, \delta > 0,$$

where $\dot{W}_{\epsilon, \delta}(t, x)$ is constructed in Section 2.2, define the Stratonovich integral of $\Psi(t, x)$ as

$$(2.12) \quad \int_{\mathbb{R}_+ \times \mathbb{R}^d} \Psi(t, x) W(dt, dx) \triangleq \lim_{\epsilon, \delta \rightarrow 0^+} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \Psi(t, x) \dot{W}_{\epsilon, \delta}(t, x) dt dx$$

whenever the limit exists in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ and is independent of the way how δ and ϵ approach to zero. We can also use the convergence in probability in above definition. But as in most works on SPDE, $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ norm is easier to deal with so that we choose the $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ convergence throughout this work. Notice that this definition implicates that $u(t, x)$ as a solution to (2.1) is in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$. After defining the Stratonovich integral, we can give the following definition about the solution.

DEFINITION 2.1. A random field $\{u(t, x), t \geq 0, x \in \mathbb{R}^d\}$ is called a mild solution to (1.1) if for any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, the random field

$$\Psi(s, y) \equiv G(t - s, x - y)u(s, y)1_{[0, t]}(s)$$

is Stratonovich integrable and if (2.1) is satisfied.

To prove Theorem 1.1, we shall use the Stratonovich expansion (see [20], [19] and references therein for the multiple Stratonovich integrals). According to the algorithm in [16], formally iterating (2.1) infinitely many times we have heuristically a solution candidate

$$(2.13) \quad u(t, x) = \sum_{n=0}^{\infty} S_n(g_n(\cdot, t, x))$$

with $S_0(g_0(\cdot, t, x)) = 1$. Here is how the notation $S_n(g_n(\cdot, t, x))$ is justified: The iteration procedure creates the recurrent relation

$$(2.14) \quad S_{n+1}(g_{n+1}(\cdot, t, x)) = \int_{\mathbb{R}^d} \int_0^t G(t - s, x - y) S_n(g_n(\cdot, s, y)) W(ds, dy).$$

Iterating this relation formally we have

$$(2.15) \quad \begin{aligned} S_n(g_n(\cdot, t, x)) &= \int_{(\mathbb{R}^d)^n} \int_{[0, t]_{<}^n} G(t - s_n, x - x_n) \cdots G(s_2 - s_1, x_2 - x_1) \\ &\quad \times W(ds_1, dx_1) \cdots W(ds_n, dx_n) \\ &= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) W(ds_1, dx_1) \cdots W(ds_n, dx_n) \quad (\text{say}), \end{aligned}$$

where $[0, t]_{<}^n := \{(s_1, \dots, s_n) \in [0, t]^n \text{ satisfies } 0 < s_1 < s_2 < \dots < s_n < t\}$, and the conventions $x_{n+1} = x$ and $s_{n+1} = t$ are adopted. Thus, the notation “ $S_n(g_n(\cdot, t, x))$ ” is reasonably introduced for an n -multiple Gaussian integral of the integrand

$$(2.16) \quad \begin{aligned} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \\ = \left(G(t - s_n, x - x_n) \cdots G(s_2 - s_1, x_2 - x_1) \right) 1_{[0, t]_{<}^n}(s_1, \dots, s_n) \end{aligned}$$

($n = 1, 2, \dots$). In Section 4, the Stratonovich integrability of $g_n(\cdot, t, x)$ shall be rigorously established in Theorem 4.1.

The above construction indicates that the existence of the system (2.1) can be implied by the convergence of the random series defined by (2.13) in an appropriate form. This will be justified rigorously in Section 5 as part of the proof of Theorem 1.1.

DEFINITION 2.2. Let $f : (\mathbb{R}_+ \times \mathbb{R}^d)^n \rightarrow \mathbb{R}$ be measurable such that for every $\epsilon, \delta > 0$

$$\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \in \mathcal{L}^2(\Omega, \mathcal{F}, \mathbb{P}).$$

Then we define the n -multiple Stratonovich integral of f as

$$(2.17) \quad \begin{aligned} S_n(f) &:= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) W(ds_1, dx_1) \cdots W(ds_n, dx_n) \\ &= \lim_{\epsilon, \delta \rightarrow 0^+} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) \\ &\quad \times ds_1 \cdots ds_n dx_1 \cdots dx_n \end{aligned}$$

whenever the limit exists $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$.

REMARK 2.3. Along with the set-up of our model, the Stratonovich integrand f is given as a generalized function for $d = 3$. Indeed ([32]), Definition 2.2 can be extended to the setting of generalized functions f . A detail is provided near the end of this section for the construction needed in $d = 3$.

The following lemma provides a convenient test of Stratonovich integrability that we shall use in this work.

LEMMA 2.4. *The n -multiple Stratonovich integral $S_n(f)$ exists if and only if the limit*

$$\lim_{\substack{(\epsilon, \delta) \rightarrow 0^+ \\ (\epsilon', \delta') \rightarrow 0^+}} \mathbb{E} \left\{ \int_{(\mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \right\} \\ \times \left\{ \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) \left(\prod_{k=1}^n \dot{W}_{\epsilon', \delta'}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \right\}$$

exists

PROOF. The existence of the limit in (2.17) is another way to say that the family

$$\mathcal{Z}_{\epsilon, \delta} = \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} f(s_1, \dots, s_n, x_1, \dots, x_n) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n$$

is a Cauchy sequence in $\mathcal{L}^2(\Omega, \mathcal{F}, \mathbb{P})$ as $\epsilon, \delta \rightarrow 0^+$, which is equivalent to the lemma. $\square$

Definition 2.2 can be extended to a random field $f(s_1, \dots, s_n, x_1, \dots, x_n)$ in an obvious way. Most of the time in this paper, however, we deal with a deterministic integrand and demand some effective ways to compute the expectation of multiple Stratonovich integral of deterministic integrands.

LEMMA 2.5. *In the setting of a deterministic integrand, the $\mathcal{L}^2$ -convergence in Definition 2.2 can be replaced $\mathcal{L}^p$ -convergence for any $p \geq 2$.*

PROOF. This has been proved in the setting of time-independent noise (Theorem 6.2 and Remark 6.3, [14]). In the context of the general integrand, one can identify $f(s_1, \dots, s_n, x_1, \dots, x_n)$ with $f(x_1, \dots, x_n)$ and the time-dependent Gaussian field $\dot{W}(t, x)$ with the time-independent Gaussian field $\dot{W}(x)$ by viewing the time variable as the extra dimension of the space variables. Then Theorem 6.2 and Remark 6.3, [14] applies to our setting. $\square$

This lemma brings some convenience in computation. An example of such is the Fubini theorem: Given the integers $l_1, \dots, l_m \geq 1$ and the l_j -multiple time-space variate functions f_j ($1 \leq j \leq m$), the Stratonovich integrabilities of $f_1, \dots, f_m$ implies the Stratonovich integrability of $f_1 \otimes \cdots \otimes f_m$ and

$$(2.18) \quad S_{l_1 + \dots + l_m}(f_1 \otimes \cdots \otimes f_m) = \prod_{j=1}^m S_{l_j}(f_j).$$

Let us recall an identity [31, p.201, Lemma 5.2.6] known as Wick's formula which states that

$$(2.19) \quad \begin{cases} \mathbb{E} \prod_{k=1}^{2n} g_k = \sum_{\mathcal{D} \in \Pi_n} \prod_{(j,k) \in \mathcal{D}} \mathbb{E} g_j g_k \\ \mathbb{E} \prod_{k=1}^{2n-1} g_k = 0, \end{cases}$$

where $(g_1, \dots, g_{2n})$ is a mean zero normal vector, and Π_n is the set of all pair partitions of $\{1, 2, \dots, 2n\}$. As a side remark, $\#(\Pi_n) = \frac{(2n)!}{2^n n!}$. Applying (2.19) to $g_k = \dot{W}_{\epsilon, \delta}(s_k, x_k)$ in the case of deterministic integrand f , and taking the (ϵ, δ) -limit, we have

$$(2.20) \quad \mathbb{E} S_{2n-1}(f) = 0$$

and

$$(2.21) \quad \begin{aligned} \mathbb{E} S_{2n}(f) &= \sum_{\mathcal{D} \in \Pi_n} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} f(s_1, \dots, s_{2n}, x_1, \dots, x_{2n}) \\ &\quad \times \left(\prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \right) ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \end{aligned}$$

under the Stratonovich integrability. In particular, the expectation of a $(2n)$ -multiple Stratonovich integral is non-negative if the integrand is non-negative. Take $f = g_{2n}(\cdot, t, 0)$ for example. In the case of integrability

$$\begin{aligned} &\mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) \\ &= \sum_{\mathcal{D} \in \Pi_n} \int_{[0, t]^{2n}} \int_{(\mathbb{R}^d)^{2n}} ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \left(G(t - s_{2n}, -x_{2n}) \cdots G(s_2 - s_1, x_2 - x_1) \right) \\ &\quad \times \left(\prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \right). \end{aligned}$$

With the substitutions $s_k \mapsto t - s_{2n-k+1}$ and $x_k \mapsto -x_{2n-k+1}$ ($k = 1, 2, \dots, 2n$) performed on the right hand side,

$$\begin{aligned} (2.22) \quad &\mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) \\ &= \sum_{\mathcal{D} \in \Pi_n} \int_{[0, t]^{2n}} \int_{(\mathbb{R}^d)^{2n}} ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ &\quad \times \left(\prod_{(j,k) \in \mathcal{D}} |s_{2n-j+1} - s_{2n-k+1}|^{-\alpha_0} \gamma(x_{2n-j+1} - x_{2n-k+1}) \right) \\ &= \sum_{\mathcal{D} \in \Pi_n} \int_{[0, t]^{2n}} \int_{(\mathbb{R}^d)^{2n}} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ &\quad \times \left(\prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \right) ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \end{aligned}$$

where we follow the convention on the right hand side that $s_0 = 0$ and $x_0 = 0$.

Since the condition (1.5) encompasses the cases where the covariance function $\gamma(\cdot)$ exists only as a generalized function (e.g., $\gamma(\cdot) = \delta_0(\cdot)$ in $d = 1$), the meaning of the multiple integrals on the right hand side of (2.21) needs to be clarified. Indeed, by (2.9) and (2.19)

$$\begin{aligned}
 (2.23) \quad & \mathbb{E} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} f(s_1, \dots, s_{2n}, x_1, \dots, x_{2n}) \left(\prod_{k=1}^{2n} \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) \\
 & \times ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \\
 & = \sum_{\mathcal{D} \in \Pi_n} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} f(s_1, \dots, s_{2n}, x_1, \dots, x_{2n}) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_\delta^0(s_j - s_k) \gamma_{2\epsilon}(x_j - x_k) \right) \\
 & \times ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n}
 \end{aligned}$$

where

$$(2.24) \quad \gamma_\delta^0(u) = \Gamma(\alpha_0)^{-1} \int_0^{\delta^{-1}} e^{-\lambda|u|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \quad \text{and} \quad \gamma_\epsilon(x) = \int_{\mathbb{R}^d} \gamma(y) p_\epsilon(x - y) dy.$$

Inspired by (2.17) and in light of (2.23), we therefore define

$$\begin{aligned}
 (2.25) \quad & \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} f(s_1, \dots, s_{2n}, x_1, \dots, x_{2n}) \left(\prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \right) \\
 & \times ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n} \\
 & \triangleq \lim_{\epsilon, \delta \rightarrow 0^+} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} f(s_1, \dots, s_{2n}, x_1, \dots, x_{2n}) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_\delta^0(s_j - s_k) \gamma_\epsilon(x_j - x_k) \right) \\
 & \times ds_1 \cdots ds_{2n} dx_1 \cdots dx_{2n}
 \end{aligned}$$

whenever the limit exists.

To end this section we take the chance to address an inconvenient fact from (2.5) where $G(t, x)$ is defined as a measure rather than a function on $\mathbb{R}^3$. In this case, the integrand $g_n(\cdot, t, x)$ introduced in (2.16) appears as an integrand of the n -multiple Stratonovich integral in the sense given below: For any $\eta > 0$, set

$$G_\eta(t, x) = \int_{\mathbb{R}^3} p(\eta, x - y) G(t, y) dy$$

where $p(t, x)$ is the Brownian semi-group on $\mathbb{R}^3$. Write

$$\begin{aligned}
 & g_{n, \eta}(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \\
 & = \left(G_\eta(t - s_n, x - x_n) \cdots G_\eta(s_2 - s_1, x_2 - x_1) \right) 1_{[0, t] <^n (s_1, \dots, s_n)}.
 \end{aligned}$$

The n -multiple Stratonovich integral is defined in the following two steps:

$$\begin{aligned}
 & \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \\
 & := \lim_{\eta \rightarrow 0^+} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_{n, \eta}(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n
 \end{aligned}$$

for any $\epsilon, \delta > 0$, and

$$\begin{aligned} S(g_n(\cdot, t, x)) &:= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) W(ds_1, x_1) \cdots W(s_n, x_n) \\ &:= \lim_{\epsilon, \delta \rightarrow 0^+} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \left(\prod_{k=1}^n \dot{W}_{\epsilon, \delta}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \end{aligned}$$

whenever both limits exist in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$. It will be verified later that the limits do exist under the assumption (1.5). Therefore, it can be treated together with dimensions $d = 1, 2$.

3. Stratonovich moment bound. Let the pair partition $\mathcal{D} \in \Pi_n$ be fixed. Let $\mu_{j,k}(d\xi)$ $((j, k) \in \mathcal{D})$ be the finite measures on $\mathbb{R}^d$ such that

$$(3.1) \quad \gamma_{j,k}(x) \equiv \int_{\mathbb{R}^d} e^{i\xi \cdot x} \mu_{j,k}(d\xi) \geq 0 \quad x \in \mathbb{R}^d.$$

Since $\mu_{j,k}$ are finite, $\gamma_{j,k}(x)$ are defined point-wise.

Let $A_{j,k} \subset \mathbb{R}_+$ $((j, k) \in \mathcal{D})$ be measurable and define

$$(3.2) \quad \gamma_{j,k}^0(r) = \Gamma(\alpha_0)^{-1} \int_{A_{j,k}} e^{-\lambda|r|} \frac{d\lambda}{\lambda^{1-\alpha_0}} \quad r \in \mathbb{R}.$$

Clearly, $\gamma_{j,k}^0(\cdot)$ is non-negative and non-negative definite (see the trick played in (2.10) for the second claim).

For any non-negative function $\mathcal{G}(t, x)$ on $\mathbb{R}_+ \times \mathbb{R}^d$, we introduce the notations

$$(3.3) \quad \|\mathcal{G}\|^{(0)} \equiv \int_{\mathbb{R}_+ \times \mathbb{R}^d} \mathcal{G}(t, x) dt dx \quad (\text{It is } (j, k)\text{-index free!}),$$

$$(3.4) \quad \|\mathcal{G}\|_{j,k}^{(1)} \equiv \Gamma(\alpha_0)^{-1} \int_{A_{j,k} \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} e^{-\lambda t + i\xi \cdot x} \mathcal{G}(t, x) dt dx \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu_{j,k}(d\xi),$$

$$(3.5) \quad \|\mathcal{G}\|_{j,k}^{(2)} \equiv \left(\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} \gamma_{j,k}^0(s-t) \gamma_{j,k}(x-y) \mathcal{G}(s, x) \mathcal{G}(t, y) ds dx dt dy \right)^{1/2}.$$

The aim of this section is to provide a meaningful bound for the multiple integral

$$\begin{aligned} &\int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2}} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2}} dx_1 \cdots dx_{2n} \\ &\times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \end{aligned}$$

with the non-negative functions $G_l(t, x)$ $(l = 1, 2, \dots, 2n)$ such that $\|G_l\|^{(0)}, \|G_l\|_{j,k}^{(1)}, \|G_l\|_{j,k}^{(2)} < \infty$, where n_1 and n_2 are non-negative integers with $2n = n_1 + n_2$, $I_1 = \{1, \dots, n_1\}$ and $I_2 = \{n_1 + 1, \dots, 2n\}$, and where the following conventions are adopted: $s_0 = 0$, $x_0 = 0$, and $s_{n_1} = 0$, $x_{n_1} = 0$ in the expression $G_{n_1+1}(s_{n_1+1} - s_{n_1}, x_{n_1+1} - x_{n_1})$. The settings $n_1 = 2n$ (and therefore $n_2 = 0$) and $n_2 = 2n$ (and therefore $n_1 = 0$) are considered as the special cases where the above integral takes the form

$$\begin{aligned} &\int_{(\mathbb{R}_+)^{2n}} ds_1 \cdots ds_{2n} \int_{\mathbb{R}^{2n}} dx_1 \cdots dx_{2n} \\ &\times \left(\prod_{l=1}^{2n} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k). \end{aligned}$$

THEOREM 3.1. *There is a partition $\{Q_0, Q_1, Q_2\}$ of $\{1, \dots, 2n\}$, possibly depending on n_1, n_2 and $\mathcal{D}$, such that $\#(Q_2)$ is even and $\#(Q_0) = \#(Q_1)$ and that*

$$(3.6) \quad \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2}} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2}} dx_1 \cdots dx_{2n} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \\ \leq \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2\|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)$$

where the subscripts $\{(j, k) \in \mathcal{D}\}$ are distributed in the Q_1 -product and in the Q_2 -product in the following way: First, the pair (j, k) appearing in the Q_1 -product does not appear in the Q_2 -product, and vice versa. Second, the pair appearing in Q_1 -product appears exactly once (so the number of different (j, k) in Q_1 -product is $\#(Q_1)$), while the pair in Q_2 -product appears exactly twice (so the number of different (j, k) in Q_2 -product is $2^{-1}\#(Q_2)$).

To further illustrate labeling system used on the right hand side of (3.6), we consider the special case $n = 2$ and $\mathcal{D} = \{(1, 2), (3, 4)\}$. A possible bound on the right hand side of (3.6) is

$$\left(\|G_1\|^{(0)} \right) \left(\|G_2\|_{1,2}^{(1)} \right) \left(\|G_3\|_{3,4}^{(2)} \|G_4\|_{3,4}^{(2)} \right).$$

where $Q_0 = \{1\}$, $Q_1 = \{2\}$ and $Q_2 = \{3, 4\}$.

In another special case $n = 3$, $\mathcal{D} = \{(1, 5), (2, 4), (3, 6)\}$, a possible bound on the right hand side of (3.6) is

$$\left(\|G_1\|^{(0)} \|G_6\|^{(0)} \right) \left(\|G_3\|_{1,5}^{(1)} \|G_5\|_{2,4}^{(1)} \right) \left(\|G_2\|_{3,6}^{(2)} \|G_4\|_{3,6}^{(2)} \right)$$

where $Q_0 = \{1, 6\}$, $Q_1 = \{3, 5\}$ and $Q_2 = \{2, 4\}$.

PROOF. We carry out argument by induction on n : When $n = 1$, there are three possible forms for the left hand: " $n_1 = n_2 = 1$ ", " $n_1 = 2$ and $n_2 = 0$ " or " $n_1 = 0$ and $n_2 = 2$ ".

When $n_1 = n_2 = 1$, the left hand of (3.6) is

$$\int_{\mathbb{R}_+^2} ds_1 ds_2 \int_{(\mathbb{R}^d)^2} dx dy G_1(s_1, x) G_2(s_2, y) \gamma_{1,2}^0(s_1 - s_2) \gamma_{1,2}(x - y) \leq \|G_1\|_{1,2}^{(2)} \|G_2\|_{1,2}^{\{2\}}$$

where the inequality follows from Cauchy-Schwartz's inequality. So the claim holds with $Q_0 = Q_1 = \emptyset$ and $Q_2 = \{1, 2\}$.

When $n_1 = 2$ and $n_2 = 0$, the left hand side is

$$\int \int_{\{s_1 < s_2\}} ds_1 ds_2 \int_{(\mathbb{R}^d)^2} dx dy G_1(s_1, x) G_2(s_2 - s_1, y - x) \gamma_{1,2}^0(s_1 - s_2) \gamma_{1,2}(x - y) \\ = \left(\int_{\mathbb{R}_+ \times \mathbb{R}^d} G_1(s_1, x) ds_1 dx \right) \left(\int_{\mathbb{R}_+ \times \mathbb{R}^d} G_2(t, y) \gamma_{1,2}^0(t) \gamma_{1,2}(y) dt dy \right).$$

Notice

$$\begin{aligned}
& \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_2(t, y) \gamma_{1,2}^0(t) \gamma_{1,2}(y) dt dy \\
&= \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_2(t, y) \left(\int_{A_{1,2} \times \mathbb{R}^d} e^{-\lambda t + i\xi \cdot y} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu_{1,2}(d\xi) \right) dt dy \\
&= \Gamma(\alpha_0)^{-1} \int_{A_{1,2} \times \mathbb{R}^d} \left(\int_{\mathbb{R}_+ \times \mathbb{R}^d} G_2(t, y) e^{-\lambda t + i\xi \cdot y} dt dy \right) \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu_{1,2}(d\xi) \leq \|G_2\|_{1,2}^{(1)}.
\end{aligned}$$

In summary

$$\begin{aligned}
& \iint_{\{s_1 < s_2\}} ds_1 ds_2 \int_{(\mathbb{R}^d)^2} dx dy G_1(s_1, x) G_2(s_2 - s_1, y - x) |s_1 - s_2|^{-\alpha_0} \gamma(x - y) \\
& \leq \|G_1\|^{(0)} \|G_2\|_{1,2}^{(1)}.
\end{aligned}$$

Thus, the claim holds with $Q_0 = \{1\}$, $Q_1 = \{2\}$ and $Q_2 = \phi$.

Similarly, when $n_1 = 0$ and $n_2 = 2$, the claim holds with the bound $\|G_1\|_{1,2}^{(1)} \|G_2\|^{(0)}$ and $Q_0 = \{2\}$, $Q_1 = \{1\}$ and $Q_2 = \phi$.

Assume the claim holds for $n - 1$. We now verified it for n . Assume that j_0 and k_0 are paired with $2n$ and n_1 , respectively, i.e. $(j_0, 2n), (k_0, n_1) \in \mathcal{D}$. The idea is to separate

$$G_{2n}(s_{2n} - s_{2n-1}, x_{2n} - x_{2n-1}) \gamma_{j_0, 2n}^0(s_{j_0} - s_{2n}) \gamma_{j_0, 2n}(x_{j_0} - x_{2n})$$

or

$$G_{n_1}(s_{n_1} - s_{n_1-1}, x_{n_1} - x_{n_1-1}) \gamma_{k_0, n_1}^0(s_{k_0} - s_{n_1}) \gamma_{k_0, n_1}(x_{k_0} - x_{n_1}),$$

whichever possible, from the multiple time-space integral.

We consider the following three possible cases: Case 1: $j_0 \in I_2$ or $k_0 \in I_1$. In other words, at least one of n_1 and $2n$ has a domestic pair. In the remaining settings, both n_1 and $2n$ have inter-group pairs. We shall treat it in the following two different cases: Case 2: $j_0 = n_1$ and $k_0 = 2n$, i.e., $(n_1, 2n) \in \mathcal{D}$; and Case 3: $1 \leq j_0 < n_1 < k_0 < 2n$.

Case 1: We actually claim a better bound

$$\begin{aligned}
(3.7) \quad & \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2}} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2}} dx_1 \cdots dx_{2n} \\
& \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \\
& \leq \frac{1}{2} \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)
\end{aligned}$$

in this case for the argument needed in Case 3.

Due to similarity, we only consider the case $j_0 \in I_2$. For $s_{j_0} \leq s_{2n-1} \leq s_{2n}$,

$$\gamma_{j_0, 2n}^0(s_{2n} - s_{j_0}) \leq \gamma_{j_0, 2n}^0(s_{2n} - s_{2n-1}).$$

Thus, the left hand side of (3.7) yields to the bound

$$\begin{aligned} & \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2-1}} ds_1 \cdots ds_{2n-1} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2-1}} dx_1 \cdots dx_{2n-1} \\ & \times \left(\prod_{\rho=1}^2 \prod_{l \in I'_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}'} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \right) \int_{s_{2n-1}}^\infty ds_{2n} \\ & \times \int_{\mathbb{R}^d} dx_{2n} G_{2n}(s_{2n} - s_{2n-1}, x_{2n} - x_{2n-1}) \gamma_{j_0,2n}^0(s_{2n} - s_{2n-1}) \gamma_{j_0,2n}(x_{2n} - x_{j_0}) \end{aligned}$$

where $I'_1 = I_1$ and $I'_2 = I_2 \setminus \{2n\} = \{n_1 + 1, \dots, 2n - 1\}$ and $\mathcal{D}' = \mathcal{D} \setminus (j_0, 2n) \in \Pi_{n-1}$. Notice

$$\begin{aligned} & \int_{s_{2n-1}}^\infty ds_{2n} \int_{\mathbb{R}^d} dx_{2n} G_{2n}(s_{2n} - s_{2n-1}, x_{2n} - x_{2n-1}) \gamma_{j_0,2n}^0(s_{2n} - s_{2n-1}) \gamma_{j_0,2n}(x_{2n} - x_{j_0}) \\ & = \int_0^\infty ds \int_{\mathbb{R}^d} dx_{2n} G_{2n}(s, x_{2n} - x_{2n-1}) \gamma_{j_0,2n}^0(s) \int_{\mathbb{R}^d} e^{i\xi \cdot (x_{2n} - x_{j_0})} \mu_{j_0,2n}(d\xi) \\ & = \Gamma(\alpha_0)^{-1} \int_{A_{j_0,2n} \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \exp\{i\xi \cdot (x_{2n-1} - x_{j_0})\} \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{2n}(s, x) e^{-\lambda s + i\xi \cdot x} ds dx \\ & \leq \Gamma(\alpha_0)^{-1} \int_{A_{j_0,2n} \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{2n}(s, x) e^{-\lambda s + i\xi \cdot x} ds dx \right| = \|G_{2n}\|_{j_0,2n}^{(1)}. \end{aligned}$$

In summary, the left hand of (3.7) yields to the bound

$$\begin{aligned} & \|G_{2n}\|_{j_0,2n}^{(1)} \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2-1}} ds_1 \cdots ds_{2n-1} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2-1}} dx_1 \cdots dx_{2n-1} \\ (3.8) \quad & \times \left(\prod_{\rho=1}^2 \prod_{l \in I'_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}'} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \right). \end{aligned}$$

Denote $\tilde{I}_1 = I_1$, $\tilde{I}_2 = I_2 \setminus \{j_0, 2n\}$. When $j_0 = 2n - 1$, the right hand side is equal to

$$\begin{aligned} & \|G_{2n}\|_{j_0,2n}^{(1)} \|G_{2n-1}\|^{(0)} \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2-2}} ds_1 \cdots ds_{2n-2} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2-2}} dx_1 \cdots dx_{2n-2} \\ & \times \left(\prod_{\rho=1}^2 \prod_{l \in I'_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}'} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \right). \end{aligned}$$

Applying the induction assumption, we yield the bound

$$\begin{aligned} & \|G_{2n}\|_{j_0,2n}^{(1)} \|G_{2n-1}\|^{(0)} \left(\prod_{l \in \tilde{Q}_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in \tilde{Q}_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in \tilde{Q}_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \\ & = \frac{1}{2} \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \end{aligned}$$

where $\tilde{Q}_0$, $\tilde{Q}_1$, $\tilde{Q}_2$ form a partition of $\{1, \dots, 2(n-1)\}$ obeying the rule described by Theorem 3.1. Setting $Q_0 = \tilde{Q}_0 \cup \{2n-1\}$, $Q_1 = \tilde{Q}_1 \cup \{2n\}$ and $Q_2 = \tilde{Q}_2$ we have proved (3.7).

We now assume $j_0 < 2n - 1$. Set

$$d\tilde{\mathbf{s}} = ds_1 \cdots ds_{j_0-1} ds_{j_0+1} \cdots d_{2n-1} \quad \text{and} \quad d\tilde{\mathbf{x}} = dx_1 \cdots dx_{j_0-1} dx_{j_0+1} \cdots d_{2n-1}.$$

The bound in (3.8) can be written as

$$\begin{aligned} & \|G_{2n}\|_{j_0, 2n}^{(1)} \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2-2}} d\tilde{\mathbf{s}} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2-2}} d\tilde{\mathbf{x}} \left(\prod_{\rho=1}^2 \prod_{l \in \tilde{I}_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \times \left(\int_{s_{j_0-1}}^{s_{j_0+1}} \int_{\mathbb{R}^d} G_{j_0}(s_{j_0} - s_{j_0-1}, x_{j_0} - x_{j_0-1}) G_{j_0+1}(s_{j_0+1} - s_{j_0}, x_{j_0+1} - x_{j_0}) dx_{j_0} ds_{j_0} \right) \\ & \times \left(\prod_{(j,k) \in \mathcal{D}'} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \right). \end{aligned}$$

Notice

$$\begin{aligned} & \int_{s_{j_0-1}}^{s_{j_0+1}} \int_{\mathbb{R}^d} G_{j_0}(s_{j_0} - s_{j_0-1}, x_{j_0} - x_{j_0-1}) G_{j_0+1}(s_{j_0+1} - s_{j_0}, x_{j_0+1} - x_{j_0}) dx_{j_0} ds_{j_0} \\ & = \tilde{G}_{j_0, j_0+1}(s_{j_0+1} - s_{j_0-1}, x_{j_0+1} - x_{j_0-1}) \end{aligned}$$

where

$$\tilde{G}_{j_0, j_0+1}(t, x) = \int_0^t \int_{\mathbb{R}^d} G_{j_0}(s, y) G_{j_0+1}(t - s, x - y) dy ds.$$

Therefore, by (3.8)

$$\begin{aligned} & \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2}} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2}} dx_1 \cdots dx_{2n} \\ & \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \\ & \leq \|G_{2n}\|_{j_0, 2n}^{(1)} \int_{(\mathbb{R}_+)^{n_1} \times (\mathbb{R}_+)^{n_2-2}} d\tilde{\mathbf{s}} \int_{(\mathbb{R}^d)^{n_1} \times (\mathbb{R}^d)^{n_2-2}} d\tilde{\mathbf{x}} \left(\prod_{\rho=1}^2 \prod_{l \in \tilde{I}_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \times \tilde{G}_{j_0, j_0+1}(s_{j_0+1} - s_{j_0-1}, x_{j_0+1} - x_{j_0-1}) \left(\prod_{(j,k) \in \mathcal{D}'} \gamma_{j,k}^0(s_j - s_k) \gamma_{j,k}(x_j - x_k) \right). \end{aligned}$$

Applying the induction assumption with the functions $(2(n-1))$ of them

$$G_1, \dots, G_{j_0-1}, \tilde{G}_{j_0, j_0+1}, G_{j_0+2}, \dots, G_{2n-1}$$

with $\mathcal{D}' = \mathcal{D} \setminus \{j_0, 2n\}$ and with $2(n-1) = n_1 + (n_2 - 2)$, we have one of the three possible bounds:

$$\begin{aligned} (3.9) \quad & \|G_{2n}\|_{j_0, 2n}^{(1)} \|\tilde{G}_{j_0, j_0+1}\|_{(\cdot, \cdot)}^{(i)} \left(\prod_{l \in \tilde{Q}^{(0)}} \|G_l\|^{(0)} \right) \left(\prod_{l \in \tilde{Q}_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in \tilde{Q}_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \\ & = \frac{1}{2} \|\tilde{G}_{j_0, j_0+1}\|_{(\cdot, \cdot)}^{(i)} \left(\prod_{l \in \tilde{Q}^{(0)}} \|G_l\|^{(0)} \right) \left(\prod_{l \in \tilde{Q}_1 \cup \{2n\}} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in \tilde{Q}_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \end{aligned}$$

with $i = 0, 1, 2$, where $\tilde{Q}_0, \tilde{Q}_1, \tilde{Q}_2$ form a partition of $\{1, \dots, 2n-1\} \setminus \{j_0, j_0+1\}$. When $i = 0$, $\|\tilde{G}_{j_0, j_0+1}\|_{(\cdot, \cdot)}^{(0)} = \|\tilde{G}_{j_0, j_0+1}\|^{(0)}$ is free of (j, k) -index. As $i = 1, 2$, $\|\tilde{G}_{j_0, j_0+1}\|^{(i)} =$

$\|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(i)}$ for some $(j_1, k_1) \in \mathcal{D}'$ (recall that $\mathcal{D}'$ is a pair partition on $\{1, \dots, 2n\} \setminus \{j_0, 2n\}$).

When $i = 0$, by induction assumption $1 + \#(\tilde{Q}_0) = \#(\tilde{Q}_1)$, $\#(\tilde{Q}_2)$ is even. Recall that all $(j, k) \in \mathcal{D}'$ have been assigned into $\tilde{Q}_1$ and $\tilde{Q}_2$ according to the statement of Theorem 3.1. In particular, the number of $(j, k) \in \mathcal{D}'$ in $\tilde{Q}_1$ -product is $\#(\tilde{Q}_1)$ and the number of $(j, k) \in \mathcal{D}'$ in $\tilde{Q}_2$ -product is $2^{-1}\#(\tilde{Q}_2)$. Further

$$\|\tilde{G}_{j_0, j_0+1}\|_0 = \|G_{j_0}\|^{(0)} \|G_{j_0+1}\|^{(0)}.$$

The bound (3.7) has been verified with $Q_0 = \tilde{Q}_0 \cap \{j_0, j_0 + 1\}$, $Q_1 = \tilde{Q}_1 \cup \{2n\}$ and $Q_2 = \tilde{Q}_2$.

When $i = 1$, $\#(\tilde{Q}_0) = 1 + \#(\tilde{Q}_1)$, $\#(\tilde{Q}_2)$ is even. Recall that $\|\tilde{G}_{j_0, j_0+1}\|^{(1)} = \|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(1)}$ for some $(j_1, k_1) \in \mathcal{D}' = \mathcal{D} \setminus \{j_0, 2n\}$. All $(j, k) \in \mathcal{D}' \setminus \{(j_1, k_1)\}$ have been assigned into $\tilde{Q}_1$ and $\tilde{Q}_2$ in a way that $\#(\tilde{Q}_1) - 1$ of them are in $\tilde{Q}_1$ -product and remaining of them (the number is $2^{-1}\#(\tilde{Q}_2)$) are in $\tilde{Q}_2$ -product (each of them appears twice).

$$\begin{aligned} \|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(1)} &= \Gamma(\alpha_0)^{-1} \int_{A_{j_1, k_1} \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{j_0}(t, x) e^{-\lambda t + i\xi \cdot x} dt dx \right| \\ &\quad \times \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{j_0+1}(t, x) e^{-\lambda t + i\xi \cdot x} dt dx \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \\ &\leq \left\{ \Gamma(\alpha_0)^{-1} \int_{A_{j_1, k_1} \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{j_0}(t, x) e^{-\lambda t + i\xi \cdot x} dt dx \right|^2 \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \right\}^{1/2} \\ &\quad \times \left\{ \Gamma(\alpha_0)^{-1} \int_{A_{j_1, k_1} \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{j_0+1}(t, x) e^{-\lambda t + i\xi \cdot x} dt dx \right|^2 \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \right\}^{1/2}. \end{aligned}$$

Notice

$$\begin{aligned} &\Gamma(\alpha_0)^{-1} \int_{A_{j_1, k_1} \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} G_{j_0}(t, x) e^{-\lambda t + i\xi \cdot x} dt dx \right|^2 \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \\ &= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} \gamma_{j_1, k_1}^0(s+t) \gamma_{j, k}(x-y) G_{j_0}(s, x) G_{j_0}(t, x) ds dt dx dy \\ &\leq (\|G_{j_0}\|_{j_1, k_1}^{(2)})^2. \end{aligned}$$

Same thing happens to G_{j_0+1} -factor. Thus,

$$\|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(1)} \leq \|G_{j_0}\|_{j_1, k_1}^{(2)} \|G_{j_0+1}\|_{j_1, k_1}^{(2)}.$$

So (3.7) has been verified with $Q_0 = \tilde{Q}_0$, $Q_1 = \tilde{Q}_1 \cup \{2n\}$ and $Q_2 = \tilde{Q}_2 \cup \{j_0, j_0 + 1\}$.

When $i = 2$, $\#(\tilde{Q}_0) = \#(\tilde{Q}_1)$, $\#(\tilde{Q}_2)$ is odd. The pairs $(j, k) \in \mathcal{D}'$ are distributed in a way that $\#(\tilde{Q}_1)$ of them are in $\tilde{Q}_1$ -product and remaining of them are in $\tilde{Q}_2$ -product. All (j, k) in $\tilde{Q}_2$ -product appears twice except (j_1, k_1) (which appears once). Consequently, the number of $(j, k) \in \mathcal{D}'$ in $\tilde{Q}_1$ is $\#(\tilde{Q}_1)$ and the number of $(j, k) \in \mathcal{D}'$ in $\tilde{Q}_2$ is $2^{-1}(\#(\tilde{Q}_2) +$

1). Further

$$\begin{aligned}
& \|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(2)} \\
&= \left\{ \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} \gamma_{j_1, k_1}^0(s-r) \gamma_{j_1, k_1}(x-y) \tilde{G}_{j_0, j_0+1}(s, x) \tilde{G}_{j_0, j_0+1}(r, y) ds dx dr dy \right\}^{1/2} \\
&= \left\{ \int_{\mathbb{R}^{d+1}} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} \tilde{G}_{j_0, j_0+1}(t, x) dx dt \right|^2 \mu_{j_1, k_1}^0(d\lambda) \mu_{j, k}(d\xi) \right\}^{1/2}
\end{aligned}$$

where $\mu_{j_1, k_1}^0(d\lambda)$ is the spectral measure of $\gamma_{j_1, k_1}^0(\cdot)$ (Recall that $\gamma_{j_1, k_1}^0(\cdot)$ is non-negative-definite).

Notice

$$\begin{aligned}
& \left| \int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} \tilde{G}_{j_0, j_0+1}(t, x) dx dt \right| \\
&= \left| \left(\int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} G_{j_0}(t, x) dx dt \right) \left(\int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} G_{j_0+1}(t, x) dx dt \right) \right| \\
&\leq \|G_{j_0}\|^{(0)} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} G_{j_0+1}(t, x) dx dt \right|.
\end{aligned}$$

Therefore,

$$\begin{aligned}
& \|\tilde{G}_{j_0, j_0+1}\|_{j_1, k_1}^{(2)} \\
&\leq \|G_{j_0}\|^{(0)} \left\{ \int_{\mathbb{R}^{d+1}} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{i\lambda t + i\xi \cdot x} G_{j_0+1}(t, x) dx dt \right|^2 \mu_{j_1, k_1}^0(d\lambda) \mu_{j_1, k_1}(d\xi) \right\}^{1/2} \\
&= \|G_{j_0}\|^{(0)} \|G_{j_0+1}\|_{j_1, k_1}^{(2)}.
\end{aligned}$$

The bound (3.7) has been varified with $Q_0 = \tilde{Q}_0 \cup \{j_0\}$, $Q_1 = \tilde{Q}_1 \cup \{2n\}$ and $Q_2 = \tilde{Q}_2 \cup \{j_0 + 1\}$.

Case 2. Write $\tilde{I}_1 = \{1, \dots, n_1 - 1\}$, $\tilde{I}_2 = \{n_1 + 1, \dots, 2n - 1\}$ and

$$d\tilde{\mathbf{s}} = ds_1 \cdots ds_{n_1-1} ds_{n_1+1} \cdots ds_{2n-1} \quad \text{and} \quad d\tilde{\mathbf{x}} = dx_1 \cdots dx_{n_1-1} dx_{n_1+1} \cdots dx_{2n-1}.$$

Since $(n_1, 2n) \in \mathcal{D}$ in this case, the left hand side of (3.6) is equal to

$$\begin{aligned}
& \int_{(\mathbb{R}_+)^{n_1-1} \times (\mathbb{R}_+)^{n_2-1}} d\tilde{\mathbf{s}} \int_{(\mathbb{R}^d)^{n_1-1} \times (\mathbb{R}^d)^{n_2-1}} d\tilde{\mathbf{x}} \\
&\times \left(\prod_{\rho=1}^2 \prod_{l \in \tilde{I}_\rho} G_l(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j, k) \in \tilde{\mathcal{D}}} \gamma_{j, k}^0(s_j - s_k) \gamma_{j, k}(x_j - x_k) \\
&\times \int_{s_{n_1-1}}^\infty \int_{s_{2n-1}}^\infty \int_{\mathbb{R}^d \times \mathbb{R}^d} ds_{n_1} ds_{2n} dx_{n_1} dx_{2n} \gamma_{n_1, 2n}^0(s_{2n} - s_{n_1}) \gamma_{n_1, 2n}(x_{2n} - x_{n_1}) \\
&\times G_{n_1}(s_{n_1} - s_{n_1-1}, x_{n_1} - x_{n_1-1}) G_{2n}(s_{2n} - s_{2n-1}, x_{2n} - x_{2n-1}).
\end{aligned}$$

where $\tilde{D} = \mathcal{D} \setminus \{n_1, 2n\}$. Notice

$$\begin{aligned} & \int_{s_{n_1-1}}^{\infty} \int_{s_{2n-1}}^{\infty} \int_{\mathbb{R}^d \times \mathbb{R}^d} ds_{n_1} ds_{2n} dx_{n_1} dx_{2n} \gamma_{n_1, 2n}^0(s_{2n} - s_{n_1}) \gamma_{n_1, 2n}(x_{2n} - x_{n_1}) \\ & \times G_{n_1}(s_{n_1} - s_{n_1-1}, x_{n_1} - x_{n_1-1}) G_{2n}(s_{2n} - s_{2n-1}, x_{2n} - x_{2n-1}) \\ & = \int_0^{\infty} \int_0^{\infty} \int_{\mathbb{R}^d \times \mathbb{R}^d} ds dr dx dy \gamma_{n_1, 2n}^0(s - r) \gamma_{n_1, 2n}(x - y) G_{n_1}(s, x - x_{n_1-1}) G_{2n}(r, y - x_{2n-1}) \\ & \leq \|G_{n_1}\|_{n_1, 2n}^{(2)} \|G_{2n}\|_{n_1, 2n}^{(2)} \end{aligned}$$

where last step follows from Cauchy-Schwartz inequality and then shift-invariance of the space variables.

By the induction assumption, we have the bound

$$\|G_{n_1}\|_{n_1, 2n}^{(2)} \|G_{2n}\|_{n_1, 2n}^{(2)} \left(\prod_{l \in \tilde{Q}_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in \tilde{Q}_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in \tilde{Q}_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)$$

where $\tilde{Q}_0, \tilde{Q}_1, \tilde{Q}_2$ form a partition of $\{1, \dots, 2n\} \setminus \{n_1, 2n\}$ with $\#(\tilde{Q}_0) = \#(\tilde{Q}_1)$ and $\#(\tilde{Q}_2)$ being even. All $(j, k) \in \mathcal{D}'$ are distributed according to the statement of Theorem 3.1. In particular, the number of (j, k) in $\tilde{Q}_1$ is $\#(\tilde{Q}_1)$ and number of (j, k) in $\tilde{Q}_2$ is $2^{-1} \#(\tilde{Q}_2)$. The bound (3.6) has been varified with $Q_0 = \tilde{Q}_0$, $Q_1 = \tilde{Q}_1$ and $Q_2 = \tilde{Q}_2 \cup \{n_1, 2n\}$.

Case 3. Recall that this is the case when $(j_0, 2n), (k_0, n_1) \in \mathcal{D}$ with $j_0 \in I_1, k_1 \in I_2$ and $j_0 \neq n_1, k_0 \neq 2n$. The idea essentially comes from the treatment in Case 1. The extra obstacle comes from the fact that there is no restrictive order between s_{j_0} and s_{2n} , nor between s_{k_0} and s_{n_1} , as $(j_0, 2n)$ and (k_0, n_1) are inter-group pairs. To fix it, we break the left hand side of (3.6) into two parts by the indicators $1_{\{s_{j_0} \leq s_{2n-1}\}}$ and $1_{\{s_{j_0} > s_{2n-1}\}}$

On $\{s_{2n-1} \geq s_{j_0}\}$, $\gamma_{j_0, 2n}^0(s_{2n} - s_{j_0}) \leq \gamma_{j_0, 2n}^0(s_{2n} - s_{2n-1})$. By a strategy same as the one used for (3.9), the first part of the decomposition yields the bound in (3.7), i.e.,

$$\frac{1}{2} \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right).$$

On $\{s_{2n-1} < s_{j_0}\}$, on the other hand, $s_{n_1-1} \geq s_{j_0} > s_{2n-1} \geq s_{k_0}$. Therefore $\gamma_{n_1, k_0}^0(s_{n_1} - s_{k_0}) \leq \gamma_{n_1, k_0}^0(s_{n_1} - s_{n_1-1})$. Repeating the same procedure (with $2n$ being replaced by n_1), the second part of the decomposition has the bound (3.7), i.e.,

$$\frac{1}{2} \left(\prod_{l \in Q'_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q'_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q'_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)$$

for the partition Q'_1, Q'_2, Q'_3 of $\{1, \dots, 2n\}$ that meets all requirement given in Theorem 3.1.

Therefore, the left hand side of (3.6) is less than or equal to

$$\begin{aligned} & \frac{1}{2} \left\{ \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \right. \\ & \left. + \left(\prod_{l \in Q'_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q'_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q'_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right) \right\}. \end{aligned}$$

Between the partitions $\{Q_0, Q_1, Q_2\}$ and $\{Q'_0, Q'_1, Q'_2\}$, choosing the one that produces the larger product completes the induction. $\square$

4. Stratonovich integrability of $g_n(\cdot, t, x)$. Let $n \geq 1$ be fixed and recall that $g_n(\cdot, t, x)$ is defined in (2.16). Through this section, we adopt the following notations:

$$\epsilon = (\epsilon_1, \dots, \epsilon_n), \quad \tilde{\epsilon} = (\epsilon_{n+1}, \dots, \epsilon_{2n}) \quad \text{and} \quad \delta = (\delta_1, \dots, \delta_n), \quad \tilde{\delta} = (\tilde{\delta}_{n+1}, \dots, \tilde{\delta}_{2n})$$

$$\bar{\epsilon} = (\epsilon, \tilde{\epsilon}) = (\epsilon_1, \dots, \epsilon_{2n}) \quad \text{and} \quad \bar{\delta} = (\delta, \tilde{\delta}) = (\delta_1, \dots, \delta_{2n})$$

for $\epsilon_1, \dots, \epsilon_{2n}, \delta_1, \dots, \delta_{2n} > 0$. The notation " $\epsilon \rightarrow 0^+$ " means $\epsilon_1, \dots, \epsilon_n \rightarrow 0^+$. The notations " $\tilde{\epsilon} \rightarrow 0^+$ ", " $\delta \rightarrow 0^+$ ", " $\tilde{\delta} \rightarrow 0^+$ ", " $\bar{\epsilon} \rightarrow 0^+$ " and " $\bar{\delta} \rightarrow 0^+$ " are used in obvious way.

Set

$$(4.1) \quad S_{n,\epsilon,\delta}(g_n(\cdot, t, x)) = \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \\ \times \left(\prod_{k=1}^n \dot{W}_{\epsilon_k, \delta_k}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n.$$

The main goal of this section is

THEOREM 4.1. *Under the condition (1.5), the limit*

$$(4.2) \quad \lim_{\epsilon, \delta \rightarrow 0^+} S_{n,\epsilon,\delta}(g_n(\cdot, t, x))$$

exists in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ for each $n \geq 1$, $t > 0$ and $x \in \mathbb{R}^d$. Consequently, $g_n(\cdot, t, x)$ is n -multiple Stratonovich integrable. Further,

$$(4.3) \quad \mathbb{E} \left[S_n(g_n(\cdot, t, x)) \right]^2 = \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]^n \times [0,t]^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k)$$

where, $I_1 = \{1, \dots, n\}$, $I_2 = \{n+1, \dots, 2n\}$, where, to simplify the notation, we follow the convention on the right hand side that $s_0 = 0$, $x_0 = 0$, $s_n = 0$, $x_n = 0$ in the expression $G(s_{n+1} - s_n, x_{n+1} - x_n)$.

Notice that under the initial condition given in (1.1) and due to the stationarity of the Gaussian noise in the space variable,

$$S_{n,\epsilon,\delta}(g_n(\cdot, t, x)) \stackrel{d}{=} S_{n,\epsilon,\delta}(g_n(\cdot, t, 0))$$

we may take $x = 0$ in the proof of Theorem 4.1. By Lemma 2.4, to establish (4.2), all we need is to prove that the limit

$$(4.4) \quad \lim_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \mathbb{E} S_{n,\epsilon,\delta}(g_n(\cdot, t, 0)) S_{n,\tilde{\epsilon},\tilde{\delta}}(g_n(\cdot, t, 0))$$

exists. By Wick's formula (2.19) and the covariance identity (2.11)

$$\begin{aligned}
& \mathbb{E} S_{n,\epsilon,\delta}(g_n(\cdot, t, 0)) S_{n,\tilde{\epsilon},\tilde{\delta}}(g_n(\cdot, t, 0)) \\
&= \sum_{\mathcal{D} \in \Pi_n} \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^{2n}} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, 0) g_n(s_{n+1}, \dots, s_{2n}, x_{n+1}, \dots, x_{2n}, t, 0) \\
&\times \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \\
&= \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]_{<}^n \times [0,t]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \left(G(t - s_n, -x_n) \cdots G(s_2 - s_1, x_2 - x_1) \right) \\
&\times \left(G(t - s_{2n}, -x_{2n}) \cdots G(s_{n+2} - s_{n+1}, x_{n+2} - x_{n+1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k)
\end{aligned}$$

where $\gamma_\delta^0(\cdot)$ and $\gamma_\epsilon(\cdot)$ are defined in (2.24).

Under the substitution $s_l \mapsto t - s_{n-l+1}$, $x_l \mapsto -x_{n-l+1}$ ($1 \leq l \leq n$) and $s_l \mapsto t - s_{2n-l+1}$, $x_l \mapsto -x_{2n-l+1}$ ($n+1 \leq l \leq 2n$), the right hand side is equal to

$$\begin{aligned}
& \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]_{<}^n \times [0,t]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\
&\times \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_{\sigma(j)} \vee \delta_{\sigma(k)}}(s_{\sigma(j)} - s_{\sigma(k)}) \gamma_{\epsilon_{\sigma(j)} + \epsilon_{\sigma(k)}}(x_{\sigma(j)} - x_{\sigma(k)})
\end{aligned}$$

where σ is the permutation on $\{1, 2, \dots, 2n\}$ such that $\sigma(l) = n - l + 1$ ($l \in I_1$) and $\sigma(l) = 2n - l + 1$ ($l \in I_2$). Since $\sigma(I_1) = I_1$ and $\sigma(I_2) = I_2$, the equality continues to be equal to

$$\begin{aligned}
& \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]_{<}^n \times [0,t]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\
&\times \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k).
\end{aligned}$$

In summary,

$$\begin{aligned}
(4.5) \quad & \mathbb{E} S_{n,\epsilon,\delta}(g_n(\cdot, t, 0)) S_{n,\tilde{\epsilon},\tilde{\delta}}(g_n(\cdot, t, 0)) \\
&= \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]_{<}^n \times [0,t]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \\
&\times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k).
\end{aligned}$$

To prove Theorem 4.1, by continuity theorem of Laplace transform, part of our strategy is to establish the existence of the limit

$$\lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}_+} dt_1 dt_2 \exp \{ -\theta_1 t_1 - \theta_2 t_2 \} \int_{[0,t_1]_{<}^n \times [0,t_2]_{<}^n} ds_1 \cdots ds_{2n} \cdots$$

to the integrals on the right hand side of (4.5) for fixed δ .

To this end, let $\theta_1, \theta_2 > 0$ be fixed but arbitrary and set

$$\begin{aligned} & \tilde{g}_\rho(s_1, \dots, s_n, x_1, \dots, x_n) \\ &= \left(\prod_{l=1}^n e^{-\theta_\rho(s_l - s_{l-1})} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) 1_{(\mathbb{R}_+)^n_{<}}(s_1, \dots, s_n) \end{aligned}$$

with $\rho = 1, 2$ and the convention $s_0 = 0$ and $x_0 = 0$. Consider the function

$$\begin{aligned} (4.6) \quad F_{\bar{\epsilon}}(t_1, \dots, t_{2n}) &\equiv \int_0^{t_1} \dots \int_0^{t_{2n}} ds_1, \dots, ds_{2n} \int_{(\mathbb{R}^d)^{2n}} dx_1 \dots dx_{2n} \\ &\times \tilde{g}_1(s_1, \dots, s_n, x_1, \dots, x_n) \tilde{g}_2(s_{n+1}, \dots, s_{2n}, x_{n+1}, \dots, x_{2n}) \\ &\times \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k). \end{aligned}$$

The constructions of $\tilde{g}_\rho(s_1, \dots, s_n, x_1, \dots, x_n)$ and $F_{\bar{\epsilon}}(t_1, \dots, t_{2n})$ serves our purpose through the relation given in (4.11) below.

LEMMA 4.2. *There is a constant C independent of $\bar{\epsilon}$ such that*

$$(4.7) \quad F_{\bar{\epsilon}}(+\infty, \dots, +\infty) \leq C < \infty.$$

Further, the distribution function (up to normalization) $F_{\bar{\epsilon}}$ weakly converges as $\bar{\epsilon} \rightarrow 0^+$.

PROOF. Let $\lambda_1, \dots, \lambda_{2n} > 0$ be fixed but arbitrary.

$$\begin{aligned} & \int_{\mathbb{R}_+^{2n}} \exp \left\{ - \sum_{l=1}^{2n} \lambda_l t_l \right\} F_{\bar{\epsilon}}(dt_1, \dots, dt_{2n}) \\ &= \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right) \int_{\mathbb{R}_+^{2n}} \exp \left\{ - \sum_{l=1}^{2n} \lambda_l t_l \right\} \\ &\times \tilde{g}_1(t_1, \dots, t_n, x_1, \dots, x_n) \tilde{g}_2(t_{n+1}, \dots, t_{2n}, x_{n+1}, \dots, x_{2n}) dt_1 \dots dt_{2n}. \end{aligned}$$

There is the main reason for the Laplace transform: The weak convergence claimed in Lemma 4.2 requires that

$$\lim_{\bar{\epsilon} \rightarrow 0^+} F_{\bar{\epsilon}}(t_1, \dots, t_{2n})$$

exists at the continuous points of $F_{\bar{\epsilon}}$. This is not a trivial matter as $F_{\bar{\epsilon}}$ is not monotonic in $\epsilon_1, \dots, \epsilon_{2n}$. On the other hand, we shall see that the proposed Laplace transform installs the needed monotonicity. In addition, the proposed Laplace transform reduces the proof of (4.7) to the bounds for the moment of the Brownian intersection local times.

By our set-up

$$\begin{aligned} & \int_{\mathbb{R}_+^{2n}} \exp \left\{ - \sum_{l=1}^{2n} \lambda_l t_l \right\} \tilde{g}_1(t_1, \dots, t_n, x_1, \dots, x_n) \tilde{g}_2(t_{n+1}, \dots, t_{2n}, x_{n+1}, \dots, x_{2n}) dt_1 \dots dt_{2n} \\ &= \int_{(\mathbb{R}_+)^n_{<} \times (\mathbb{R}_+)^n_{<}} dt_1 \dots dt_{2n} \exp \left\{ - \sum_{l=1}^{2n} \lambda_l t_l \right\} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} e^{-\theta_\rho(t_l - t_{l-1})} G(t_l - t_{l-1}, x_l - x_{l-1}) \right). \end{aligned}$$

Write

$$\sum_{l=1}^{2n} \lambda_l t_l = \sum_{\rho=1}^2 \sum_{l \in I_\rho} \lambda_l t_l = \sum_{\rho=1}^2 \sum_{l \in I_\rho} c_l (t_l - t_{l-1})$$

where

$$c_l = \sum_{k=l}^n \lambda_k \quad (1 \leq l \leq n) \quad \text{and} \quad c_l = \sum_{k=n+l}^{2n} \lambda_k \quad (n+1 \leq l \leq 2n).$$

The right hand side is equal to

$$\begin{aligned} & \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} e^{-(c_l + \theta_\rho)(t_l - t_{l-1})} G(t_l - t_{l-1}, x_l - x_{l-1}) \right) \\ &= \prod_{\rho=1}^2 \prod_{l \in I_\rho} \int_0^\infty dt e^{-(c_l + \theta_\rho)t} G(t, x_l - x_{l-1}) \\ &= \left(\frac{1}{2}\right)^{2n} \prod_{\rho=1}^2 \prod_{l \in I_\rho} \int_0^\infty dt \exp \left\{ -\frac{1}{2}(c_l + \theta_\rho)^2 t \right\} p(t, x_l - x_{l-1}) \\ &= \left(\frac{1}{2}\right)^{2n} \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \\ &\quad \times \prod_{\rho=1}^2 \prod_{l \in I_\rho} \exp \left\{ -\frac{1}{2}(c_l + \theta_\rho)^2 (t_l - t_{l-1}) \right\} p(t_l - t_{l-1}, x_l - x_{l-1}) \end{aligned}$$

where $p(t, x)$ is the Brownian semi-group defined in (1.20) and the second step follows from the identity (1.19).

In summary

$$\begin{aligned} & \int_{\mathbb{R}_+^{2n}} \exp \left\{ -\sum_{l=1}^{2n} \lambda_l t_l \right\} F_\epsilon(dt_1, \dots, dt_{2n}) \\ &= \left(\frac{1}{2}\right)^{2n} \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2} \sum_{\rho=1}^2 \sum_{l \in I_\rho} (c_l + \theta_\rho)^2 (t_l - t_{l-1}) \right\} \\ &\quad \times \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} p(t_l - t_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right). \end{aligned}$$

Notice that the function

$$f(x_1, \dots, x_{2n}) = \prod_{\rho=1}^2 \prod_{l \in I_\rho} p(t_l - t_{l-1}, x_l - x_{l-1})$$

is the density of the random vector $(B_1(t_1), \dots, B_1(t_n), B_2(t_{n+1}), \dots, B_2(t_{2n}))$, where $B_1(t)$ and $B_2(t)$ are two independent d -dimensional Brownian motions. Let " $\mathbb{E}_0$ " denote the expectation of the Brownian motions and let the map $v: \{1, \dots, 2n\} \longrightarrow \{1, 2\}$ be given

by $v(I_\rho) = \{\rho\}$ ($\rho = 1, 2$). We have

$$\begin{aligned}
& \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} p(t_l - t_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right) \\
&= \mathbb{E}_0 \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\
&= \int_{(\mathbb{R}^d)^n} \exp \left\{ -\frac{1}{2} \sum_{(j,k) \in \mathcal{D}} (\epsilon_j + \epsilon_k) |\xi_{j,k}|^2 \right\} \mathbb{E}_0 \exp \left\{ i \sum_{(j,k) \in \mathcal{D}} \xi_{j,k} \cdot (B_{v(j)}(t_j) - B_{v(k)}(t_k)) \right\} \\
&\quad \times \prod_{(j,k) \in \mathcal{D}} \mu(d\xi_{j,k}) \\
&= \int_{(\mathbb{R}^d)^n} \exp \left\{ -\sum_{(j,k) \in \mathcal{D}} (\epsilon_j + \epsilon_k) |\xi_{j,k}|^2 \right\} \exp \left\{ -\frac{1}{2} \text{Var} \left(\sum_{(j,k) \in \mathcal{D}} \xi_{j,k} \cdot (B_{v(j)}(t_j) - B_{v(k)}(t_k)) \right) \right\} \\
&\quad \times \prod_{(j,k) \in \mathcal{D}} \mu(d\xi_{j,k}).
\end{aligned}$$

The right hand side is monotone in $\epsilon_1, \dots, \epsilon_{2n}$ ¹. Using monotone convergence we conclude that the limit

$$\begin{aligned}
(4.8) \quad & \lim_{\bar{\epsilon} \rightarrow 0^+} \int_{\mathbb{R}_+^{2n}} \exp \left\{ -\sum_{l=1}^{2n} \lambda_l t_l \right\} F_{\bar{\epsilon}}(dt_1, \dots, dt_{2n}) \\
&= \left(\frac{1}{2} \right)^{2n} \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2} \sum_{\rho=1}^2 \sum_{l \in I_\rho} (c_l + \theta_\rho)^2 (t_l - t_{l-1}) \right\} \\
&\quad \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k))
\end{aligned}$$

exists as a extended real number. Further,

$$\begin{aligned}
& \int_{\mathbb{R}_+^{2n}} \exp \left\{ -\sum_{l=1}^{2n} \lambda_l t_l \right\} F_{\bar{\epsilon}}(dt_1, \dots, dt_{2n}) \\
&\leq \left(\frac{1}{2} \right)^{2n} \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2} \sum_{\rho=1}^2 \sum_{l \in I_\rho} (c_l + \theta_\rho)^2 (t_l - t_{l-1}) \right\} \\
&\quad \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\
&\leq \left(\frac{1}{2} \right)^{2n} \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2} \sum_{\rho=1}^2 \sum_{l \in I_\rho} \theta_\rho^2 (t_l - t_{l-1}) \right\} \\
&\quad \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k))
\end{aligned}$$

¹This is the main reason that we introduce the Laplacian parameters $\lambda_1, \dots, \lambda_{2n}$

Noticing that

$$\sum_{\rho=1}^2 \sum_{l \in I_\rho} \theta_\rho^2(t_l - t_{l-1}) = \theta_1^2 t_n + \theta_2^2 t_{2n}$$

and taking $\lambda_1 = \dots = \lambda_{2n} = 0$ on the left hand side,

$$\begin{aligned} & F_{\bar{\epsilon}}(+\infty, \dots, +\infty) \\ & \leq \left(\frac{1}{2}\right)^{2n} \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2}(\theta_1^2 t_n + \theta_2^2 t_{2n}) \right\} \\ & \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)). \end{aligned}$$

To complete the proof, therefore, all we need is

$$\begin{aligned} (4.9) \quad & \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2}(\theta_1^2 t_n + \theta_2^2 t_{2n}) \right\} \\ & \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) < \infty. \end{aligned}$$

By the fact that

$$\theta_1 \theta_2 \int_{t_n}^{\infty} ds_1 \int_{t_{2n}}^{\infty} ds_2 \exp \left\{ -\frac{1}{2}(\theta_1^2(s_1 - t_n) + \theta_2^2(s_2 - t_{2n})) \right\} = 1.$$

and by Fubini's theorem,

$$\begin{aligned} (4.10) \quad & \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2}(\theta_1^2 t_n + \theta_2^2 t_{2n}) \right\} \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\ & = \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2}(\theta_1^2 t_n + \theta_2^2 t_{2n}) \right\} \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\ & \times \theta_1 \theta_2 \int_{t_n}^{\infty} ds_1 \int_{t_{2n}}^{\infty} ds_2 \exp \left\{ -\frac{1}{2}(\theta_1^2(s_1 - t_n) + \theta_2^2(s_2 - t_{2n})) \right\} \\ & = \theta_1 \theta_2 \int_0^{\infty} \int_0^{\infty} ds_1 ds_2 \exp \left\{ -\frac{1}{2}(\theta_1^2 s_1 + \theta_2^2 s_2) \right\} \int_{[0,s_1]^n \times [0,s_2]^n} dt_1 \cdots dt_{2n} \\ & \times \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)). \end{aligned}$$

In addition,

$$\begin{aligned} & \int_{[0,s_1]^n \times [0,s_2]^n} dt_1 \cdots dt_{2n} \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\ & \leq \int_{[0,s_1]^n \times [0,s_2]^n} dt_1 \cdots dt_{2n} \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\ & = \prod_{(j,k) \in \mathcal{D}} \int_0^{s_{v(j)}} \int_0^{s_{v(k)}} \gamma(B_{v(j)}(t_1) - B_{v(k)}(t_2)) dt_1 dt_2 \end{aligned}$$

where the last step follows from Fubini's theorem. For the mutual intersection local times on the right hand side (i.e., for the $(j, k) \in \mathcal{D}$ but j and k coming from different groups) we treat them by Fourier transform

$$\begin{aligned}
& \int_0^{s_1} \int_0^{s_2} \gamma(B_1(t_1) - B_2(t_2)) dt_1 dt_2 \\
&= \int_{\mathbb{R}^d} \mu(d\xi) \int_0^{s_1} \int_0^{s_2} \exp \left\{ i\xi \cdot (B_1(t_1) - B_2(t_2)) \right\} dt_1 dt_2 \\
&= \int_{\mathbb{R}^d} \mu(d\xi) \left[\int_0^{s_1} e^{i\xi \cdot B_1(t)} dt \right] \overline{\left[\int_0^{s_2} e^{i\xi \cdot B_2(t)} dt \right]} \\
&\leq \left\{ \int_{\mathbb{R}^d} \mu(d\xi) \left| \int_0^{s_1} e^{i\xi \cdot B_1(t)} dt \right|^2 \right\}^{1/2} \left\{ \int_{\mathbb{R}^d} \mu(d\xi) \left| \int_0^{s_2} e^{i\xi \cdot B_2(t)} dt \right|^2 \right\}^{1/2} \\
&= \left\{ \int_0^{s_1} \int_0^{s_1} \gamma(B_1(t_1) - B_1(t_2)) dt_1 dt_2 \right\}^{1/2} \left\{ \int_0^{s_2} \int_0^{s_2} \gamma(B_2(t_1) - B_2(t_2)) dt_1 dt_2 \right\}^{1/2}.
\end{aligned}$$

We therefore have

$$\begin{aligned}
& \prod_{(j,k) \in \mathcal{D}} \int_0^{s_{v(j)}} \int_0^{s_{v(k)}} \gamma(B_{v(j)}(t_1) - B_{v(k)}(t_2)) dt_1 dt_2 \\
&\leq \left[\int_0^{s_1} \int_0^{s_1} \gamma(B_1(t_1) - B_1(t_2)) dt_1 dt_2 \right]^n \left[\int_0^{s_2} \int_0^{s_2} \gamma(B_2(t_1) - B_2(t_2)) dt_1 dt_2 \right]^n.
\end{aligned}$$

Summarizing the steps since (4.10),

$$\begin{aligned}
& \mathbb{E}_0 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} dt_1 \cdots dt_{2n} \exp \left\{ -\frac{1}{2}(\theta_1^2 t_n + \theta_2^2 t_{2n}) \right\} \prod_{(j,k) \in \mathcal{D}} \gamma(B_{v(j)}(t_j) - B_{v(k)}(t_k)) \\
&\leq \theta_1 \theta_2 \left\{ \int_0^\infty \exp \left\{ -\frac{\theta_1^2}{2} s \right\} \mathbb{E}_0 \left[\int_0^s \int_0^s \gamma(B(t_1) - B(t_2)) dt_1 dt_2 \right]^n ds \right\} \\
&\times \left\{ \int_0^\infty \exp \left\{ -\frac{\theta_2^2}{2} s \right\} \mathbb{E}_0 \left[\int_0^s \int_0^s \gamma(B(t_1) - B(t_2)) dt_1 dt_2 \right]^n ds \right\} \\
&< \infty
\end{aligned}$$

where the last step follows from (6.1), Lemma 6.1 in [14]. $\square$

LEMMA 4.3. *For any $\mathcal{D} \in \Pi_n$, $t_1, t_2 > 0$ and $\eta > 0$, the limit*

$$\begin{aligned}
& \lim_{\bar{\epsilon} \rightarrow 0^+} \int_{[0, t_1]_{\leq}^n \times [0, t_2]_{\leq}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\
& \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_\eta^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k)
\end{aligned}$$

exists and is finite.

PROOF. The weak convergence of F_ϵ^- established in Lemma 4.2 can be stated as the existence of the limit

$$\lim_{\bar{\epsilon} \rightarrow 0^+} \int_{\mathbb{R}_+^{2n}} \varphi(t_1, \dots, t_{2n}) F_\epsilon^-(dt_1, \dots, dt_{2n})$$

for any bounded, continuous and non-negative function $\varphi(t_1, \dots, t_{2n})$ on $\mathbb{R}_+^{2n}$. On the other hand, similar to (4.10),

(4.11)

$$\begin{aligned}
& \int_{\mathbb{R}_+^{2n}} \varphi(t_1, \dots, t_{2n}) F_{\bar{\epsilon}}(dt_1, \dots, dt_{2n}) \\
&= \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} e^{-\theta_\rho(s_l - s_{l-1})} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\
&\quad \times \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \\
&= (\theta_1 \theta_2) \int_{\mathbb{R}_+^2} dt_1 dt_2 \exp\{-\theta_1 t_1 - \theta_2 t_2\} \int_{[0, t_1]_{\mathbb{Z}}^n \times [0, t_2]_{\mathbb{Z}}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\
&\quad \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k).
\end{aligned}$$

We reach the conclusion that the limit

$$\begin{aligned}
& \lim_{\bar{\epsilon} \rightarrow 0^+} \int_{\mathbb{R}_+^2} dt_1 dt_2 \exp\{-\theta_1 t_1 - \theta_2 t_2\} \int_{[0, t_1]_{\mathbb{Z}}^n \times [0, t_2]_{\mathbb{Z}}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\
&\quad \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k)
\end{aligned}$$

exists for any $\theta_1, \theta_2 > 0$. Set

$$\begin{aligned}
\mathcal{G}_{\bar{\epsilon}}^{\mathcal{D}}(t_1, t_2) &= \int_{[0, t_1]_{\mathbb{Z}}^n \times [0, t_2]_{\mathbb{Z}}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\
&\quad \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k).
\end{aligned}$$

By continuity theorem of Laplace transform again, $\mathcal{G}_{\bar{\epsilon}}^{\mathcal{D}}(t_1, t_2)$ weakly converges: There is a non-decreasing and right continuous function $\mathcal{G}^{\mathcal{D}}(t_1, t_2)$ such that

$$(4.12) \quad \lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}}^{\mathcal{D}}(t_1, t_2) = \mathcal{G}^{\mathcal{D}}(t_1, t_2)$$

at every continuous point (t_1, t_2) of $\mathcal{G}^{\mathcal{D}}(\cdot, \cdot)$. (Actually, Theorem 5.22, [27] is stated for probability measures on $(\mathbb{R}^+)^d$. The case of general measures on $(\mathbb{R}^+)^d$ can be derived as in the proof of Theorem 2a, Section 1, Chapter 2, [24] Although this theorem only considers measures on $\mathbb{R}_+$ its extension to $\mathbb{R}_+^2$ is routine).

We now claim that $\mathcal{G}^{\mathcal{D}}(\cdot, \cdot)$ is continuous on $\mathbb{R}_+ \times \mathbb{R}_+$. Consequently, this is to say that (4.12) holds on $\mathbb{R}_+ \times \mathbb{R}_+$ and therefore Lemma 4.3 holds. To prove it, all we need is to show that

$$\lim_{\eta_1, \eta_2 \rightarrow 0^+} \sup_{\bar{\epsilon}} \left\{ \mathcal{G}_{\bar{\epsilon}}^{\mathcal{D}}(t_1, t_2) - \mathcal{G}_{\bar{\epsilon}}^{\mathcal{D}}(t_1 - \eta_1, t_2 - \eta_2) \right\} = 0$$

for every $(t_1, t_2) \in \mathbb{R}_+ \times \mathbb{R}_+$, Or,

$$\lim_{\eta_1, \eta_2 \rightarrow 0^+} \sup_{\bar{\epsilon}} \int_{([0, t_1]_{<}^n \times [0, t_2]_{<}^n) \setminus ([0, t_1 - \eta_1]_{<}^n \times [0, t_2 - \eta_2]_{<}^n)} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) = 0.$$

Indeed,

$$\int_{([0, t_1]_{<}^n \times [0, t_2]_{<}^n) \setminus ([0, t_1 - \eta_1]_{<}^n \times [0, t_2 - \eta_2]_{<}^n)} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \varphi(s_1, \dots, s_{2n}) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \\ \leq \sup_{t_1, \dots, t_{2n}} \varphi(t_1, \dots, t_{2n}) \int_{([0, t_1]_{<}^n \times [0, t_2]_{<}^n) \setminus ([0, t_1 - \eta_1]_{<}^n \times [0, t_2 - \eta_2]_{<}^n)} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k).$$

Thus, our claim follows from the fact (established in the proof of Lemma 3.6, [14]) that

$$\lim_{\eta_1, \eta_2 \rightarrow 0^+} \sup_{\bar{\epsilon}} \int_{([0, t_1]_{<}^n \times [0, t_2]_{<}^n) \setminus ([0, t_1 - \eta_1]_{<}^n \times [0, t_2 - \eta_2]_{<}^n)} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) = 0$$

Thus, (4.12) holds for every $(t_1, t_2) \in \mathbb{R}_+ \times \mathbb{R}_+$. Taking

$$\varphi(t_1, \dots, t_{2n}) = \prod_{(j,k) \in \mathcal{D}} \gamma_\eta^0(t_j - t_k)$$

in (4.12) completes the proof. $\square$

LEMMA 4.4. *For any $\mathcal{D} \in \Pi_n$, $t_1, t_2 > 0$, the limit*

$$\lim_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \int_{[0, t_1]_{<}^n \times [0, t_2]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k)$$

exists and finite.

REMARK 4.5. In view (4.5), Lemma 4.4 leads to the existence of the limit in (4.4), and therefore (Lemma 2.4) to the existence of the $\mathcal{L}^2$ -limit in (4.2). According to the definition given in (2.25), Lemma 4.4 justifies the use of the notation

$$\int_{[0, t_1]_{<}^n \times [0, t_2]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k).$$

Therefore, taking limit in (4.5) leads to (4.3). That completes the proof of Theorem 4.1

PROOF. Let

$$\begin{aligned} \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) &= \int_{[0, t_1]_{<}^n \times [0, t_2]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ &\times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k). \end{aligned}$$

For each $\eta > 0$

$$\mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) = \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \Big|_{\delta_1 = \cdots = \delta_{2n} = \eta}$$

and

$$\begin{aligned} \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2) &= \int_{[0, t_1]_{<}^n \times [0, t_2]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ &\times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k). \end{aligned}$$

By the monotonicity of $\mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2)$ in $\delta_1 \cdots \delta_{2n}$ and by (1.21),

$$\mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) \leq \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \leq \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2)$$

whenever $\delta_1, \dots, \delta_{2n} < \eta$. Therefore,

$$\lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) \leq \liminf_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \leq \liminf_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \leq \limsup_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2)$$

where the limit on the left end is guaranteed by Lemma 4.3. To complete the proof, all we need is that

$$(4.13) \quad \lim_{\eta \rightarrow 0^+} \left\{ \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2) - \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) \right\} = 0 \quad \text{uniformly over } \bar{\epsilon}.$$

Indeed, (4.13) implies that for any $\eta_1 > 0$,

$$\limsup_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2) \leq \eta_1 + \lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2)$$

as η is sufficiently small. Since $\lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) < \infty$ according to Lemma 4.3, in particular, $\limsup_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2) < \infty$. On the other hand, by the fact that $\lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2)$ is non-increasing in η , the limit

$$\lim_{\eta \rightarrow 0^+} \lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2)$$

exists and is finite (as it is bounded by $\limsup_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, 0}(t_1, t_2)$). So we have

$$\lim_{\eta \rightarrow 0^+} \lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2) \leq \liminf_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \leq \liminf_{\bar{\epsilon}, \bar{\delta} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \bar{\delta}}(t_1, t_2) \leq \eta_1 + \lim_{\eta \rightarrow 0^+} \lim_{\bar{\epsilon} \rightarrow 0^+} \mathcal{G}_{\bar{\epsilon}, \eta}(t_1, t_2)$$

Letting $\eta_1 \rightarrow 0^+$ leads to the proof of Lemma 4.4.

It remains to prove (4.13). For being consistent to the notation $\gamma_\delta^0(\cdot)$ for time-covariance, by (1.21) we use $\gamma_0^0(\cdot)$ instead of $|\cdot|^{-\alpha_0}$. Notice

$$\begin{aligned} & \prod_{(j,k) \in \mathcal{D}} \gamma_0^0(s_j - s_k) - \prod_{(j,k) \in \mathcal{D}} \gamma_\eta^0(s_j - s_k) \\ & \leq \sum_{(j_1, k_1) \in \mathcal{D}} (\gamma_0^0 - \gamma_\eta^0)(s_{j_1} - s_{k_1}) \prod_{(j,k) \in \mathcal{D} \setminus \{(j_1, k_1)\}} \gamma_0^0(s_j - s_k). \end{aligned}$$

All we need is to show that for every $(j_1, k_1) \in \mathcal{D}$

$$\begin{aligned} & \int_{[0, t_1]_{\leq}^n \times [0, t_2]_{\leq}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \times \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right) \left((\gamma_0^0 - \gamma_\eta^0)(s_{j_1} - s_{k_1}) \prod_{(j,k) \in \mathcal{D} \setminus \{(j_1, k_1)\}} \gamma_0^0(s_j - s_k) \right) \end{aligned}$$

converges to 0 uniformly over $\bar{\epsilon}$.

We are in the position of using Theorem 3.1. Set

$$\gamma_{j_1, k_1}^0(u) = (\gamma_0^0 - \gamma_\eta^0)(\cdot) = \Gamma(\alpha_0)^{-1} \int_{\eta^{-1}}^{\infty} e^{-\lambda|u|} \frac{d\lambda}{\lambda^{1-\alpha_0}}$$

and $\gamma_{j,k}^0(\cdot) = \gamma_0^0(u)$ for $(j,k) \neq (k_1, j_1)$. The above integral is written as

$$\begin{aligned} & \int_{[0, t_1]_{\leq}^n \times [0, t_2]_{\leq}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \times \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right) \\ & \leq e^{t_1 + t_2} \int_{(\mathbb{R}_+)^2} \exp\{-\tilde{t}_1 - \tilde{t}_2\} \int_{[0, \tilde{t}_1]_{<}^n \times [0, \tilde{t}_2]_{\leq}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ & \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right) \\ & = e^{t_1 + t_2} \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^{2n}} d\mathbf{x} \\ & \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} e^{-(s_l - s_{l-1})} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma_{j,k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \right). \end{aligned}$$

Applying Theorem 3.1 to $G_l(t, x) = e^{-t} G(t, x)$ ($1 \leq l \leq 2n$), the integral on the right hand side yields the bound

$$\left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)$$

where Q_0, Q_1, Q_2 form a partition of $\{1, \dots, 2n\}$ with $\#(Q_0) = \#(Q_1)$ and $\#(Q_2)$ is even. What important to our course is that (j_1, k_1) either appears in Q_1 -product once or in Q_2 twice. By (2.4)

$$\|G_l\|^{(0)} = \int_0^\infty \int_{\mathbb{R}^d} e^{-t} G(t, x) dx dt = \int_0^\infty t e^{-t} dt = 1 \quad l = 1, 2, \dots, 2n.$$

The rest of the argument is to check that $\|G_l\|_{j,k}^{(i)}$ ($i = 1, 2$) are bounded uniformly over $\bar{\epsilon}$ for $(j, k) \neq (j_1, k_1)$ and $\|G_l\|_{j_1, k_1}^{(i)} \rightarrow 0^+$ ($i = 1, 2$) uniformly over $\bar{\epsilon}$ as $\eta \rightarrow 0^+$.

For $(j, k) \neq (j_1, k_1)$,

$$\begin{aligned} \|G_l\|_{j,k}^{(1)} &= \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{-(\lambda+1)t+i\xi \cdot x} G(t, x) dx dt \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \exp \left\{ -(\epsilon_j + \epsilon_k) |\xi|^2 \right\} \mu(d\xi) \\ &\leq \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{-(\lambda+1)t+i\xi \cdot x} G(t, x) dx dt \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \end{aligned}$$

and

(4.14)

$$\begin{aligned} \|G_l\|_{j,k}^{(2)} &= \left(\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma_{\epsilon_j + \epsilon_k}(x-y) e^{-s} G(s, x) e^{-t} G(t, x) ds dx dt dy \right)^{1/2} \\ &\leq \left(\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma(x-y) e^{-s} G(s, x) e^{-t} G(t, x) ds dx dt dy \right)^{1/2} \end{aligned}$$

where the inequality follows from the procedure of Fourier transform

$$\begin{aligned} &\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma_{\epsilon_j + \epsilon_k}(x-y) e^{-s} G(s, x) e^{-t} G(t, x) ds dx dt dy \\ &= \int_{\mathbb{R}^{d+1}} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \exp \left\{ -t + i\lambda t + i\xi \cdot x \right\} G(t, x) dt dx \right|^2 \mu^0(d\lambda) \exp \left\{ -(\epsilon_j + \epsilon_k) |\xi|^2 \right\} \mu(d\xi) \\ &\leq \int_{\mathbb{R}^{d+1}} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \exp \left\{ -t + i\lambda t + i\xi \cdot x \right\} G(t, x) dt dx \right|^2 \mu^0(d\lambda) \mu(d\xi) \\ &= \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma(x-y) e^{-s} G(s, x) e^{-t} G(t, x) ds dx dt dy \end{aligned}$$

where $\mu^0(d\lambda)$ is the spectral measure of $|\cdot|^{-\alpha_0}$.

By Lemma A.1 (with $\theta = 1$), all above bounds are finite.

As for $(j, k) = (j_1, k_1)$,

$$\|G_l\|_{j_1, k_1}^{(1)} \leq \Gamma(\alpha_0)^{-1} \int_{\eta^{-1}}^\infty d\lambda \int_{\mathbb{R}^d} \frac{1}{(\lambda+1)^2 + |\xi|^2} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \rightarrow 0 \quad (\eta \rightarrow 0^+),$$

$$\begin{aligned} \|G_l\|_{j_1, k_1}^{(2)} &= \left\{ \int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} \left(\gamma_0^0 - \gamma_\eta^0 \right) (s-t) \gamma_{\epsilon_j + \epsilon_k}(x-y) e^{-t} G(t, x) e^{-s} G(s, y) ds dx dt dy \right\}^{1/2} \\ &\leq \left\{ \int_{\mathbb{R}_+ \times \mathbb{R}_+} ds dt e^{-(s+t)} \left(\gamma_0^0 - \gamma_\eta^0 \right) (s-t) \int_{\mathbb{R}^d \times \mathbb{R}^d} \gamma(x-y) G(t, x) G(s, y) dx dy \right\}^{1/2} \end{aligned}$$

where the inequality follows for the reason similar to the one for (4.14). By (A.2), Lemma A.1 (with $\theta = 1$) and dominating control theorem, the right hand side converges as $\eta \rightarrow 0^+$.

□

5. Proof of Theorem 1.1. For the part (1) of Theorem 1.1, we prove that the Stratonovich expansion (2.13) solves (2.1). According to Definition 2.1, all we need is to show

- (i) The random series in (2.13) converges in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$.
- (ii) The random field $\Psi(s, y) = G(t - s, x - y)u(s, y)1_{[0, t]}(s)$ is Stratonovich integrable and satisfies (2.1).

Our way for (i) is to show

$$(5.1) \quad \sum_{n=0}^{\infty} \left\{ \mathbb{E} [S_n(g_n(\cdot, t, x))]^2 \right\}^{1/2} < \infty \quad \forall t > 0.$$

To establish (ii), the argument given below requires a slightly stronger form of Stratonovich integrability than the one given in Definition 2.2. That is, the single limiting parameters $\epsilon > 0$ and $\delta > 0$ in (2.17) are replaced by the multi-dimensional indexes

$$\epsilon = (\epsilon_1, \dots, \epsilon_n) = (\epsilon_{n+1}, \dots, \epsilon_{2n}) = \tilde{\epsilon} \quad \text{and} \quad \delta = (\delta_1, \dots, \delta_n) = (\delta_{n+1}, \dots, \delta_{2n}) = \tilde{\delta}$$

(see (4.5)). To establish the bounds requested by (5.1), our first step is to bound the smoothed Stratonovich moments $\mathbb{E} [S_{n, \epsilon, \delta}(g_n(\cdot, t, 0))]^2$ (see (5.3) and (5.4) below). By (4.5), we have

$$(5.2) \quad \begin{aligned} & \int_0^\infty \int_0^\infty d\bar{t}_1 d\bar{t}_2 \exp \left\{ -\frac{n}{t}(\bar{t}_1 + \bar{t}_2) \right\} \mathbb{E} [S_{n, \epsilon, \delta}(g_n(\cdot, \bar{t}_1, 0)) S_{n, \epsilon, \delta}(g_n(\cdot, \bar{t}_2, 0))] \\ &= \sum_{\mathcal{D} \in \Pi_n} \int_0^\infty \int_0^\infty d\bar{t}_1 d\bar{t}_2 \exp \left\{ -\frac{n}{t}(\bar{t}_1 + \bar{t}_2) \right\} \int_{[0, \bar{t}_1]_{\mathbb{Z}} \times [0, \bar{t}_2]_{\mathbb{Z}}} ds_1 \cdots ds_{2n} \\ & \quad \times \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \quad \times \prod_{(j, k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k) \\ &= \sum_{\mathcal{D} \in \Pi_n} \left(\frac{t}{n} \right)^2 \int_{(\mathbb{R}_+)^n \times (\mathbb{R}_+)^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ & \quad \times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} e^{-nt^{-1}(s_l - s_{l-1})} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \quad \times \prod_{(j, k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j + \epsilon_k}(x_j - x_k). \end{aligned}$$

According to Theorem 3.1, each term on the right hand side yields a bound of the form

$$\left(\frac{t}{n} \right)^2 \left(\prod_{l \in Q_0} \|G_l\|^{(0)} \right) \left(\prod_{l \in Q_1} 2 \|G_l\|_{(\cdot, \cdot)}^{(1)} \right) \left(\prod_{l \in Q_2} \|G_l\|_{(\cdot, \cdot)}^{(2)} \right)$$

for $G_l(\bar{t}, x) = e^{-nt^{-1}\bar{t}} G(\bar{t}, x)$ ($l = 1, \dots, 2n$). Further

$$\|G_l\|^{(0)} = \int_0^\infty \int_{\mathbb{R}^d} e^{-nt^{-1}\bar{t}} G(\bar{t}, x) dx d\bar{t} = \int_0^\infty e^{-nt^{-1}\bar{t}} \bar{t} d\bar{t} = \left(\frac{t}{n} \right)^{2n}$$

where the first equality follows from (2.4). By (i), Lemma A.1

$$\begin{aligned} \|G_l\|_{j, k}^{(1)} &= \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{-(nt^{-1} + \lambda)\bar{t} + i\xi \cdot x} G(\bar{t}, x) dx d\bar{t} \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \exp\{-(\epsilon_j + \epsilon_k)|\xi|^2\} \mu(d\xi) \\ &\leq \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{-(nt^{-1} + \lambda)\bar{t} + i\xi \cdot x} G(\bar{t}, x) dx d\bar{t} \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \leq C < \infty. \end{aligned}$$

By (ii), Lemma A.1 with $\theta = nt^{-1}$,

$$\begin{aligned} \|G_l\|_{j,k}^{(2)} &= \left(\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma_{\epsilon_j+\epsilon_k}(x-y) e^{-nt^{-1}\bar{s}} G(s,x) e^{-nt^{-1}\bar{t}} G(\bar{t},x) ds dx dt dy \right)^{1/2} \\ &\leq \left(\int_{(\mathbb{R}_+ \times \mathbb{R}^d)^2} |s-t|^{-\alpha_0} \gamma(x-y) e^{-nt^{-1}\bar{s}} G(s,x) e^{-nt^{-1}\bar{t}} G(\bar{t},x) ds dx dt dy \right)^{1/2} \leq C \frac{t}{n} \end{aligned}$$

where the second step follows for the same reason as the one in (4.14). Therefore, the bound can be rewritten as

$$\left(\frac{t}{n}\right)^2 \left(\frac{t}{n}\right)^{2\#(Q_0)} (2C)^{\#(Q_1)} \left(\frac{t}{n}\right)^{\#(Q_2)} \leq C^n \left(\frac{t}{n}\right)^2 \left(\frac{t}{n}\right)^{2(\#(Q_0)+2^{-1}\#(Q_2))}.$$

According to Theorem 3.1, $\#(Q_0) + \#(Q_1) + \#(Q_2) = 2n$ and $\#(Q_0) = \#(Q_1)$. Therefore, $\#(Q_0) + 2^{-1}\#(Q_2) = n$. In summary,

$$\begin{aligned} &\int_0^\infty \int_0^\infty d\bar{t}_1 d\bar{t}_2 \exp\left\{-\frac{n}{t}(\bar{t}_1 + \bar{t}_2)\right\} \int_{[0,\bar{t}_1]_{<}^n \times [0,\bar{t}_2]_{<}^n} ds_1 \cdots ds_{2n} \int_{(\mathbb{R}^d)^n \times (\mathbb{R}^d)^n} dx_1 \cdots dx_{2n} \\ &\times \left(\prod_{\rho=1}^2 \prod_{l \in I_\rho} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} \gamma_{\delta_j \vee \delta_k}^0(s_j - s_k) \gamma_{\epsilon_j+\epsilon_k}(x_j - x_k) \\ &\leq C^n \left(\frac{t}{n}\right)^{2n+2} \end{aligned}$$

where the constant $C > 0$ is independent of ϵ , n , t and $\mathcal{D}$. By (5.2), therefore,

$$\int_0^\infty d\bar{t} \exp\left\{-\frac{n}{t}\bar{t}\right\} \mathbb{E}[S_{n,\epsilon,\delta}(g_n(\cdot, \bar{t}, 0))]^2 \leq \#(\Pi_n) C^n \left(\frac{t}{n}\right)^{2n+2} = C^n \frac{(2n)!}{2^n n!} \left(\frac{t}{n}\right)^{2n+2}.$$

On the other hand,

$$\begin{aligned} &\int_0^\infty \int_0^\infty d\bar{t}_1 d\bar{t}_2 \exp\left\{-\frac{n}{t}(\bar{t}_1 + \bar{t}_2)\right\} \mathbb{E}[S_{n,\epsilon,\delta}(g_n(\cdot, \bar{t}_1, 0)) S_{n,\epsilon,\delta}(g_n(\cdot, \bar{t}_2, 0))] \\ &\geq \mathbb{E}[S_{n,\epsilon,\delta}(g_n(\cdot, 0))]^2 \int_t^\infty \int_0^\infty d\bar{t}_1 d\bar{t}_2 d\bar{t}_1 d\bar{t}_2 \exp\left\{-\frac{n}{t}(\bar{t}_1 + \bar{t}_2)\right\} \\ &= \left(\frac{t}{n}\right)^2 e^{-2n} \mathbb{E}[S_{n,\epsilon,\delta}(g_n(\cdot, t, 0))]^2. \end{aligned}$$

In summary, we have the bound

$$(5.3) \quad \mathbb{E}[S_{n,\epsilon,\delta}(g_n(\cdot, t, 0))]^2 \leq C^n \frac{(2n)!}{2^n n!} \left(\frac{t}{n}\right)^{2n} \leq C^n \frac{t^{2n}}{n!}$$

with the constant $C > 0$ independent of $\epsilon = (\epsilon_1, \dots, \epsilon_n)$, $\delta = (\delta_1, \dots, \delta_n)$, n and t . Letting $\epsilon, \delta \rightarrow 0^+$. By (4.2), therefore,

$$(5.4) \quad \mathbb{E}[S_n(g_n(\cdot, t, 0))]^2 \leq C^n \frac{t^{2n}}{n!}.$$

In particular, we have (5.1).

To confirm (ii). Let $\epsilon_1, \delta_1 > 0$. In view of (2.13),

$$\begin{aligned} (5.5) \quad &1 + \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) u(s, y) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds \\ &= 1 + \sum_{n=1}^\infty \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds. \end{aligned}$$

By taking the limit properly in (5.3)

$$\mathbb{E} \left[\int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds \right]^2 \leq \frac{C^n}{n!} t^{2n} \quad n = 0, 1, \dots$$

In view of the definition in (2.12), by dominated convergence theorem, all we need is

(5.6)

$$\lim_{\epsilon_1, \delta_1 \rightarrow 0^+} \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds = S_n(g_n(\cdot, t, x))$$

in $\mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P})$ for each $n \geq 1$. Indeed, let $\epsilon' = (\epsilon_2, \dots, \epsilon_n)$ and $\delta' = (\delta_2, \dots, \delta_n)$. By the definition of $g_n(\cdot, t, x)$ given in (2.16),

$$\begin{aligned} & \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1, \epsilon', \delta'}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds \\ &= \int_{(\mathbb{R}^+ \times \mathbb{R}^d)^n} g_n(s_1, \dots, s_n, x_1, \dots, x_n, t, x) \left(\prod_{k=1}^n W_{\epsilon_k, \delta_k}(s_k, x_k) \right) ds_1 \cdots ds_n dx_1 \cdots dx_n \\ &= S_{n, \epsilon, \delta}(g_n(\cdot, t, x)). \end{aligned}$$

Therefore, by (4.2) in Theorem 4.1,

$$\begin{aligned} & \lim_{\epsilon_1, \delta_1 \rightarrow 0^+} \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds \\ &= \lim_{\epsilon_1, \delta_1 \rightarrow 0^+} \lim_{\epsilon', \delta' \rightarrow 0^+} \int_0^t \int_{\mathbb{R}^d} G(t-s, y-x) S_{n-1, \epsilon', \delta'}(g_{n-1}(\cdot, s, y)) \dot{W}_{\epsilon_1, \delta_1}(s, y) dy ds \\ &= \lim_{\epsilon, \delta \rightarrow 0^+} S_{n, \epsilon, \delta}(g_n(\cdot, t, x)) = S_n(g_n(\cdot, t, x)) \quad \text{in } \mathcal{L}^2(\Omega, \mathcal{A}, \mathbb{P}). \end{aligned}$$

So we have proved (5.6), and therefore Part (1) of Theorem 1.1.

To prove Part (2) of Theorem 1.1, all we need is to show that the condition (1.5) is necessary for

$$\mathbb{E} S_2(g_2(\cdot, t, 0)) < \infty$$

with any $t > 0$. Indeed, by (2.22)

$$\begin{aligned} & \mathbb{E} S_2(g_2(\cdot, t, 0)) \\ &= \int_{[0, t]_{<}^2} ds_1 ds_2 \int_{(\mathbb{R}^d)^2} (s_2 - s_1)^{-\alpha_0} \gamma(x_2 - x_1) G(s_1, x_1) G(s_2 - s_1, x_2 - x_1) dx_1 dx_2 \\ &= \int_{[0, t]_{<}^2} ds_1 ds_2 (s_2 - s_1)^{-\alpha_0} \int_{(\mathbb{R}^d)^2} \gamma(x_2 - x_1) G(s_1, x_1) G(s_2 - s_1, x_2 - x_1) dx_1 dx_2 \\ &= \int_{[0, t]_{<}^2} (s_2 - s_1)^{-\alpha_0} \left(\int_{\mathbb{R}^d} G(s_1, x) dx \right) \left(\int_{\mathbb{R}^d} \gamma(x) G(s_2 - s_1, x) dx \right) ds_1 ds_2 \\ &= \int_{[0, t]_{<}^2} s_1 (s_2 - s_1)^{-\alpha_0} \left[\int_{\mathbb{R}^d} \frac{\sin(|\xi|(s_2 - s_1))}{|\xi|} \mu(d\xi) \right] ds_1 ds_2 \\ &= \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{|\xi|} \int_0^t s_1 \left[\int_0^{t-s_1} \frac{\sin(s_2 |\xi|)}{s_2^{\alpha_0}} ds_2 \right] ds_1 \\ &= \int_{\mathbb{R}^d} \frac{\mu(d\xi)}{|\xi|^{2-\alpha_0}} \int_0^t s_1 \left[\int_0^{(t-s_1)|\xi|} \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 \right] ds_1. \end{aligned}$$

Notice that

$$(5.7) \quad \int_0^a \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 > 0$$

for any $0 < a \leq \pi$. When $a > \pi$,

$$\int_0^a \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 \geq \int_0^\pi \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 + \int_\pi^{2\pi} \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 \equiv \delta > 0.$$

In particular, (5.7) holds for any $a > 0$. Therefore,

$$\begin{aligned} & \mathbb{E} S_2(g_2(\cdot, t, 0)) \\ & \geq \int_{\{|\xi| \geq 2\pi t^{-1}\}} \frac{\mu(d\xi)}{|\xi|^{2-\alpha_0}} \int_0^{t/2} s_1 \left(\int_0^{(t-s_1)|\xi|} \frac{\sin s_2}{s_2^{\alpha_0}} ds_2 \right) ds_1 \\ & \geq \delta \int_{\{|\xi| \geq 2\pi t^{-1}\}} \frac{\mu(d\xi)}{|\xi|^{2-\alpha_0}} \int_0^{t/2} s_1 ds_1 = \delta \frac{t^2}{8} \int_{\{|\xi| \geq 2\pi t^{-1}\}} \frac{\mu(d\xi)}{|\xi|^{2-\alpha_0}}. \end{aligned}$$

Clearly, the finiteness on the left hand side leads to the condition (1.5). $\square$

REMARK 5.1. From (2.13) and (4.4), we have the bound

$$(5.8) \quad \mathbb{E} u^2(t, x) \leq e^{Ct^2}$$

for large t .

6. A Stratonovich moment representation. Let $\beta(t)$ and $B(t)$ be 1-dimensional and d -dimensional Brownian motions, respectively. In the rest of the paper, we assume the independence among $\beta(t)$, $B(t)$ and $\dot{W}(t, x)$ and use the notations $\mathbb{E}_0$ and $\mathbb{P}_0$ for the expectation and probability with respect to $\beta(t)$ and $B(t)$ when $B(0) = 0$ and $\beta(0) = 0$ (which is the case for most of the time).

THEOREM 6.1. Under the assumption (1.5),

$$\begin{aligned} (6.1) \quad & \int_0^\infty e^{-\theta t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt \\ & = \frac{\theta}{2} \left(\frac{1}{2}\right)^{3n} \frac{1}{n!} \int_0^\infty dt \exp\left\{-\frac{\theta^2}{2}t\right\} \\ & \quad \times \mathbb{E}_0 \left[\int_0^t \int_0^t \left((\theta(s-r) + i(\beta(s) - \beta(r))) \operatorname{sgn}(s-r) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \end{aligned}$$

for any $\theta > 0$ and $n \geq 1$.

REMARK 6.2. The appearance of the Brownian motion $\beta(t)$ is responsible for the difference in local behaviors between hyperbolic and parabolic equations when it comes to the setting of time-dependent Gaussian field. By the well-known Feynman-Kac formula, $\mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0))$ in the parabolic case is given as a constant multiple of the form

$$\frac{C^n}{n!} \mathbb{E}_0 \left[\int_0^t \int_0^t |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n$$

where the time-singularity contributed by the Gaussian field is measured by $|s - r|^{-\alpha_0}$ for close r and s . In the hyperbolic system, the time-singularity brought by the Gaussian field is measured by

$$\left| |s - r| + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \approx \left| (s - r)^2 + |s - r| \right|^{-\alpha_0/2} \approx |s - r|^{-\alpha_0/2}$$

for close r and s . It explains, for example, how the gap between conditions (1.5) and (1.7) of existence is created.

PROOF. Since $\gamma(\cdot)$ may exist as generalized function, we may encounter some legality issue. Thanks to Theorem 4.1, we are allowed to proceed with a point-wise defined $\gamma(\cdot)$, for otherwise we use $\gamma_\epsilon(\cdot)$ instead.

Our starting point is (2.22). Let $\mathcal{D} \in \Pi_n$ be fixed. For any $(j, k) \in \mathcal{D}$, we introduce the rule that $j < k$. Consequently, by (1.21)

$$\prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} = (\Gamma(\alpha_0))^{-n} \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \exp \left\{ - \sum_{(j,k) \in \mathcal{D}} \lambda_{j,k} (s_k - s_j) \right\}$$

as $(s_1, \dots, s_{2n}) \in [0, t]_{<}^{2n}$. Write

$$\sum_{(j,k) \in \mathcal{D}} \lambda_{j,k} (s_k - s_j) = \sum_{l=1}^{2n} q_l s_l = \sum_{l=1}^{2n} \left(\sum_{i=l}^{2n} q_i \right) (s_l - s_{l-1}) = \sum_{l=1}^{2n} c_l (s_l - s_{l-1})$$

where q_l is equal to $\lambda_{j,k}$ or $-\lambda_{j,k}$ for some $(j, k) \in \mathcal{D}$. Since $j < k$,

$$c_l = \sum_{i=l}^{2n} q_i = \sum_{j < l \leq k} \lambda_{j,k} \geq 0 \quad 1 \leq l \leq 2n.$$

We have

$$\begin{aligned} & \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \\ &= (\Gamma(\alpha_0))^{-n} \int_{\mathbb{R}_+^n} \prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \prod_{l=1}^{2n} e^{-c_l (s_l - s_{l-1})} G(s_l - s_{l-1}, x_l - x_{l-1}). \end{aligned}$$

Using the identity (Lemma 2.2.7, p.39, [5])

$$(6.2) \quad \theta \int_0^\infty e^{-\theta t} \int_{[0,t]_{<}^{2n}} \left(\prod_{l=1}^{2n} \varphi_l(s_l - s_{l-1}) \right) ds_1 \cdots ds_{2n} = \prod_{l=1}^{2n} \int_0^\infty \varphi_l(t) dt$$

we have

$$\begin{aligned} & \int_0^\infty dt e^{-\theta t} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \\ &= (\Gamma(\alpha_0))^{-n} \theta^{-1} \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \prod_{l=1}^{2n} \int_0^\infty e^{-\theta t} e^{-c_l t} G(t, x_l - x_{l-1}) dt. \end{aligned}$$

Noticing $c_l \geq 0$, by (1.19)

$$\begin{aligned} \int_0^\infty e^{-\theta t} e^{-c_l t} G(t, x_l - x_{l-1}) dt &= \int_0^\infty e^{-(\theta+c_l)t} G(t, x_l - x_{l-1}) dt \\ &= \frac{1}{2} \int_0^\infty \exp \left\{ -\frac{1}{2}(\theta + c_l)^2 t \right\} p(t, x_l - x_{l-1}) dt \\ &= \frac{1}{2} \int_0^\infty \exp \left\{ -\frac{\theta^2}{2} t \right\} \exp \left\{ -\theta c_l t - \frac{c_l^2}{2} t \right\} p(t, x_l - x_{l-1}) dt \end{aligned}$$

where $p(t, x)$ is the Brownian semi-group introduced in (1.20). Thus

$$\begin{aligned} &\int_0^\infty dt e^{-\theta t} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \\ &= (\Gamma(\alpha_0))^{-n} \left(\frac{1}{2} \right)^{2n} \frac{1}{\theta} \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \\ &\quad \times \prod_{l=1}^{2n} \int_0^\infty \exp \left\{ -\frac{\theta^2}{2} t \right\} \exp \left\{ -\theta c_l t - \frac{c_l^2}{2} t \right\} p(t, x_l - x_{l-1}) dt \\ &= (\Gamma(\alpha_0))^{-n} \left(\frac{1}{2} \right)^{2n} \frac{\theta}{2} \int_0^\infty dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \\ &\quad \times \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \prod_{l=1}^{2n} \exp \left\{ -\theta c_l (s_l - s_{l-1}) - \frac{c_l^2}{2} (s_l - s_{l-1}) \right\} p(s_l - s_{l-1}, x_l - x_{l-1}) \end{aligned}$$

where the last step follows from (6.2).

Multiplying the factor

$$\prod_{(j,k) \in \mathcal{D}} \gamma(x_j - x_k)$$

and integrating over $(x_1, \dots, x_{2n})$ on the both sides,

$$\begin{aligned} &\int_0^\infty dt e^{-\theta t} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ &\quad \times \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \\ &= (\Gamma(\alpha_0))^{-n} \left(\frac{1}{2} \right)^{2n} \frac{\theta}{2} \int_0^\infty dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \\ &\quad \times \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \left(\prod_{l=1}^{2n} \exp \left\{ -\theta c_l (s_l - s_{l-1}) - \frac{c_l^2}{2} (s_l - s_{l-1}) \right\} \right) \\ &\quad \times \int_{(\mathbb{R}^d)^{2n}} \left(\prod_{l=1}^{2n} p(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma(x_j - x_k) \right) dx_1 \cdots dx_{2n}. \end{aligned}$$

By the fact that

$$f(x_1, \dots, x_{2n}) = \prod_{l=1}^{2n} p(s_l - s_{l-1}, x_l - x_{l-1})$$

is the joint density of the random vector $(B(s_1), \dots, B(s_{2n}))$,

$$\begin{aligned} & \int_{(\mathbb{R}^d)^{2n}} \left(\prod_{l=1}^{2n} p(s_l - s_{l-1}, x_l - x_{l-1}) \right) \left(\prod_{(j,k) \in \mathcal{D}} \gamma(x_j - x_k) \right) dx_1 \cdots x_{2n} \\ &= \mathbb{E}_0 \prod_{(j,k) \in \mathcal{D}} \gamma(B(s_j) - B(s_k)). \end{aligned}$$

In summary,

$$\begin{aligned} & \int_0^\infty dt e^{-\theta t} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\ & \times \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \\ &= (\Gamma(\alpha_0))^{-n} \left(\frac{1}{2} \right)^{2n} \frac{\theta}{2} \int_0^\infty dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \\ & \times \mathbb{E}_0 \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{(j,k) \in \mathcal{D}} \gamma(B(s_j) - B(s_k)) \right) \\ & \times \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \exp \left\{ -\theta \sum_{l=1}^{2n} c_l(s_l - s_{l-1}) - \frac{1}{2} \sum_{l=1}^{2n} c_l^2(s_l - s_{l-1}) \right\}. \end{aligned}$$

Notice that

$$\begin{aligned} & \exp \left\{ -\theta \sum_{l=1}^{2n} c_l(s_l - s_{l-1}) - \frac{1}{2} \sum_{l=1}^{2n} c_l^2(s_l - s_{l-1}) \right\} \\ &= \mathbb{E}_0 \exp \left\{ -\theta \sum_{l=1}^{2n} c_l(s_l - s_{l-1}) - i \sum_{l=1}^{2n} c_l(\beta(s_l) - \beta(s_{l-1})) \right\}. \end{aligned}$$

Recall that

$$\sum_{l=1}^{2n} c_l(s_l - s_{l-1}) = \sum_{(j,k) \in \mathcal{D}} \lambda_{j,k}(s_k - s_j).$$

The same algebra leads to

$$\sum_{l=1}^{2n} c_l(\beta(s_l) - \beta(s_{l-1})) = \sum_{(j,k) \in \mathcal{D}} \lambda_{j,k}(\beta(s_k) - \beta(s_j)).$$

By Fubini's theorem

$$\begin{aligned}
& \int_{\mathbb{R}_+^n} \left(\prod_{(j,k) \in \mathcal{D}} \frac{d\lambda_{j,k}}{\lambda_{j,k}^{1-\alpha_0}} \right) \exp \left\{ -\theta \sum_{l=1}^{2n} c_l (s_l - s_{l-1}) - \frac{1}{2} \sum_{l=1}^{2n} c_l^2 (s_l - s_{l-1}) \right\} \\
&= \mathbb{E}_0 \prod_{(j,k) \in \mathcal{D}} \int_{\mathbb{R}_+} \exp \left\{ -\theta \lambda (s_k - s_j) - i\lambda (\beta(s_k) - \beta(s_j)) \right\} \frac{d\lambda}{\lambda^{1-\alpha_0}} \\
&= \Gamma(\alpha_0)^n \mathbb{E}_0 \prod_{(j,k) \in \mathcal{D}} \left(\theta (s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right)^{-\alpha_0}
\end{aligned}$$

where the last step follows from the identity (p. 183, [30])

$$(6.3) \quad (u + iv)^{-\alpha_0} = \Gamma(\alpha_0)^{-1} \int_0^\infty e^{-\lambda(u+iv)} \frac{d\lambda}{\lambda^{1-\alpha_0}} \quad (u, v) \in \mathbb{R}_+ \times \mathbb{R}$$

which appears to be a complex extension of (1.21).

Summarizing our steps,

$$\begin{aligned}
& \int_0^\infty dt e^{-\theta t} \int_{(\mathbb{R}^d)^{2n}} dx_1 \cdots dx_{2n} \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{l=1}^{2n} G(s_l - s_{l-1}, x_l - x_{l-1}) \right) \\
& \times \prod_{(j,k) \in \mathcal{D}} |s_j - s_k|^{-\alpha_0} \gamma(x_j - x_k) \\
&= \left(\frac{1}{2} \right)^{2n} \frac{\theta}{2} \int_0^\infty dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \\
& \times \mathbb{E}_0 \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{(j,k) \in \mathcal{D}} \left(\theta (s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right)^{-\alpha_0} \gamma(B(s_k) - B(s_j)) \right).
\end{aligned}$$

Summing up over $\mathcal{D} \in \Pi_n$, by (2.22)

$$\begin{aligned}
& \int_0^\infty e^{-\theta t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt = \left(\frac{1}{2} \right)^{2n} \frac{\theta}{2} \int_0^\infty dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \sum_{\mathcal{D} \in \Pi_n} \\
& \times \mathbb{E}_0 \int_{[0,t]_{<}^{2n}} ds_1 \cdots ds_{2n} \left(\prod_{(j,k) \in \mathcal{D}} \left(\theta (s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right)^{-\alpha_0} \gamma(B(s_k) - B(s_j)) \right).
\end{aligned}$$

Set

$$\gamma^0(z) = \left(z \cdot \operatorname{sgn}(\Re(z)) \right)^{-\alpha_0} \quad z \in \mathbb{C}$$

Then $\gamma^0(\cdot)$ is the analytic extension of the time-covariance $|\cdot|^{-\alpha_0}$ on $\{z \in \mathbb{C}; \Re(z) \neq 0\}$ with the symmetry $\gamma^0(-z) = \gamma^0(z)$. Notice that

$$\begin{aligned}
& \sum_{\mathcal{D} \in \Pi_n} \prod_{(j,k) \in \mathcal{D}} \gamma^0 \left(\theta (s_{\sigma(k)} - s_{\sigma(j)}) + i(\beta(s_{\sigma(k)}) - \beta(s_{\sigma(j)})) \right) \gamma(B(s_{\sigma(k)}) - B(s_{\sigma(j)})) \\
&= \sum_{\mathcal{D} \in \Pi_n} \prod_{(j,k) \in \mathcal{D}} \gamma^0 \left(\theta (s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right) \gamma(B(s_k) - B(s_j))
\end{aligned}$$

for any permutation σ on $\{1, \dots, 2n\}$. Consequently,

$$\begin{aligned}
& \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]^{2n}} ds_1 \cdots ds_{2n} \prod_{(j,k) \in \mathcal{D}} \gamma^0 \left(\theta(s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right) \gamma(B(s_k) - B(s_j)) \\
&= \frac{1}{(2n)!} \sum_{\mathcal{D} \in \Pi_n} \int_{[0,t]^{2n}} ds_1 \cdots ds_{2n} \prod_{(j,k) \in \mathcal{D}} \gamma^0 \left(\theta(s_k - s_j) + i(\beta(s_k) - \beta(s_j)) \right) \gamma(B(s_k) - B(s_j)) \\
&= \frac{1}{(2n)!} \sum_{\mathcal{D} \in \Pi_n} \prod_{(j,k) \in \mathcal{D}} \int_0^t \int_0^t \left(\theta(s - r) + i(\beta(s) - \beta(r)) \right) \gamma(B(s) - B(r)) ds dr \\
&= \frac{1}{(2n)!} \#(\Pi_n) \left[\int_0^t \int_0^t \gamma^0 \left(\theta(s - r) + i(\beta(s) - \beta(r)) \right) \gamma(B(s) - B(r)) ds dr \right]^n
\end{aligned}$$

Therefore, (6.1) follows from the fact that

$$\#(\Pi_n) = \frac{(2n)!}{2^n n!}.$$

□

REMARK 6.3. The monotonic order $s_1 \leq s_2 \leq \dots \leq s_{2n}$ in the representation (2.22) is a key factor that the proof of (6.1) can get through. That is the major reason why the current idea can not work for $\mathbb{E}u^p(t, x)$ for $p > 1$.

7. The time-randomized intersection local time. We assume the assumption in Theorem 1.2, i.e, (1.8) with $\alpha_0 + \alpha < 2$ (along with other conditions required for $\gamma(\cdot)$ to be non-negative definite). Motivated by Theorem 6.1 and by the relation

$$\begin{aligned}
(7.1) \quad & \left| \int_0^t \int_0^t \left(((s - r) + i(\beta(s) - \beta(r)) \operatorname{sgn}(s - r)) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right| \\
& \leq \int_0^t \int_0^t \left| ((s - r) + i(\beta(s) - \beta(r)) \operatorname{sgn}(s - r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \\
& = \int_0^t \int_0^t \left| \theta(s - r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr
\end{aligned}$$

the main goal in this section is to establish the precise large t asymptotics for the Hamiltonian on the right hand side. The fact that the two components $(s - r)$ and $\beta(s) - \beta(r)$ have different scaling rates destroys the homogeneity of the Hamiltonian. It also suggests that the contributions from $(s - r)$ and $\beta(s) - \beta(r)$ are not equal. Very likely, one of them completely dominates the game. The puzzle we face is to tell which one of $s - r$ and $\beta(s) - \beta(r)$ is the major player. To put all possible cards on the table we start with some existing results in literature. First (Theorem 1.1, [10]), under the Dalang's condition (1.17) (or (1.8) with $\alpha < 2$),

$$(7.2) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \gamma(B(s) - B(r)) ds dr \right\} = b^{\frac{2}{2-\alpha}} \mathcal{H} \quad b > 0$$

where

$$\mathcal{H} = \sup_{g \in \mathcal{F}_d} \left\{ \int_{\mathbb{R}^d \times \mathbb{R}^d} \gamma(x - y) g^2(x) g^2(y) dx dy - \frac{1}{2} \int_{\mathbb{R}^d} |\nabla g(x)|^2 dx \right\}$$

and $\mathcal{F}_d = \{g \in W^{1,2}(\mathbb{R}^d); \|g\|_2 = 1\}$.

Applying this result to the augmented Brownian motion $\tilde{B}(t) = (\beta(t), B(t))$ and to the augmented space covariance $\tilde{\gamma}(u, x) = |u|^{-\alpha_0} \gamma(x)$ (so (1.8) holds with $\tilde{\alpha} = \alpha_0 + \alpha < 2$),

$$(7.3) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} = b^{\frac{2}{2-\alpha-\alpha_0}} \tilde{\mathcal{H}}$$

for every $b > 0$. This outlines a possible scenario of " $(\beta(s) - \beta(r))$ -domination".

The scenario of " $(s - r)$ -domination" follows from the pattern (Theorem 1.1, [13] or, (4.10), [8]) that

$$(7.4) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t |s - r|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} \\ = b^{\frac{2}{2-\alpha}} \mathcal{E}$$

for every $b > 0$, where $\mathcal{E}$ is the variation given in Theorem 1.2.

The result (7.4) can not directly apply to our setting as (7.4) requires the more restrictive assumption " $2\alpha_0 + \alpha < 2$ ", for otherwise the left hand side of (7.4) blows up even before the limit is taken. Here we point out that the proof of (7.4) ([13], [8]) is based on the relation

$$\frac{1}{t^{1-\alpha_0}} \int_0^t \int_0^t |s - r|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr = \frac{1}{t} \int_0^t \int_0^t \left| \frac{s - r}{t} \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr$$

and on an argument similar to the one used for (7.2). By a parallel (and easier) modification of this idea² that for any non-negative definite function $\tilde{\gamma}^0(\cdot) \geq 0$ on $\mathbb{R}$ such that

$$(7.5) \quad \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \tilde{\gamma}^0\left(\frac{s-r}{t}\right) \gamma(B(s) - B(r)) ds dr \right\} < \infty \quad b, t > 0$$

we have

$$(7.6) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \tilde{\gamma}^0\left(\frac{s-r}{t}\right) \gamma(B(s) - B(r)) ds dr \right\} = b^{\frac{2}{2-\alpha}} \tilde{\mathcal{E}} \quad b > 0$$

where

$$\tilde{\mathcal{E}} = \sup_{g \in \mathcal{A}_d} \left\{ \int_0^1 \int_0^1 \int_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{\gamma}^0(s-r) \gamma(x-y) g^2(s, x) g^2(r, y) dx dy ds dr \right. \\ \left. - \frac{1}{2} \int_0^1 \int_{\mathbb{R}^d} |\nabla g(x)|^2 dx ds \right\}.$$

Taking $\tilde{\gamma}^0(\cdot) = \gamma_\delta^0(\cdot)$ (introduced in (2.24)), in particular,

$$(7.7) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \gamma_\delta^0\left(\frac{s-r}{t}\right) \gamma(B(s) - B(r)) ds dr \right\} = b^{\frac{2}{2-\alpha}} \mathcal{E}_\delta \quad b > 0$$

for every $\delta > 0$, where

$$\mathcal{E}_\delta = \sup_{g \in \mathcal{A}_d} \left\{ \int_0^1 \int_0^1 \int_{\mathbb{R}^d \times \mathbb{R}^d} \gamma_\delta^0(s-r) \gamma(x-y) g^2(s, x) g^2(r, y) dx dy ds dr \right. \\ \left. - \frac{1}{2} \int_0^1 \int_{\mathbb{R}^d} |\nabla g(x)|^2 dx ds \right\}.$$

²Indeed, we refer an interested reader to the argument used Section 4.2, [8] or, to the proof of (6.16) in [9] for the proof of the upper bound; and to Section 4 and 5 in [13] for the proof of lower and upper bounds, respectively.

THEOREM 7.1. Under $\alpha_0 + \alpha < 2$,

$$(7.8) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ \left. \times \gamma(B(s) - B(r)) dr ds \right\} = b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}$$

for every $b, \theta, \eta > 0$.

REMARK 7.2. (1) Theorem 7.1 clearly highlights the pattern of (7.4) rather than (7.3).

It is sharply contrary to the local behavior described in Remark 6.2.

(2) A challenge for the proof: Playing the game of (7.4) without the ticket “ $2\alpha_0 + \alpha < 2$ ” for that game.

Proof of the lower bound of (7.8). Given $\delta > 0$,

$$\frac{1}{t^{1-\alpha_0}} \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \geq \frac{\theta^{-\alpha_0}}{t} \gamma_\delta^0 \left(\frac{|s-r| + \eta\theta^{-1}|\beta(s) - \beta(r)|}{t} \right).$$

Given $a > 0$ and on $\{\max_{s \leq t} |\beta(s)| \leq a\}$, by the definition of $\gamma_\delta^0(\cdot)$ (see (2.24))

$$\begin{aligned} & \gamma_\delta^0 \left(\frac{|s-r| + \eta\theta^{-1}|\beta(s) - \beta(r)|}{t} \right) \\ & \geq \exp \left\{ -\frac{2\theta^{-1}\eta a}{\delta t} \right\} \Gamma(\alpha_0)^{-1} \int_0^{\delta^{-1}} \exp \left\{ -\lambda \frac{|s-r|}{t} \right\} \frac{d\lambda}{\lambda^{1-\alpha_0}} \\ & = \exp \left\{ -\frac{2\theta^{-1}\eta a}{\delta t} \right\} \gamma_\delta^0 \left(\frac{s-r}{t} \right). \end{aligned}$$

So we have

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ & \geq \mathbb{E}_0 \exp \left\{ \frac{b\theta^{-\alpha_0}}{t} \exp \left\{ -\frac{2\theta^{-1}\eta a}{\delta t} \right\} \int_0^t \int_0^t \gamma_\delta^0 \left(\frac{s-r}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \mathbb{P}_0 \left\{ \max_{s \leq t} |\beta(s)| \leq a \right\} \end{aligned}$$

where the last step follows from the independence between $\beta(\cdot)$ and $B(\cdot)$.

From (7.7) (with b being replaced by $b\theta^{-\alpha_0}$)

$$\begin{aligned} & \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b\theta^{-\alpha_0}}{t} \exp \left\{ -\frac{2\theta^{-1}\eta a}{\delta t} \right\} \int_0^t \int_0^t \gamma_\delta^0 \left(\frac{s-r}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \\ & \geq b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}_\delta. \end{aligned}$$

Recall a well-known fact (e.g., (1.3), [29]) that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_0 \left\{ \max_{s \leq t} |\beta(s)| \leq a \right\} = -\frac{\pi^2}{8a^2}.$$

In summary

$$\begin{aligned} & \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\epsilon(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ & \geq b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}_\delta - \frac{\pi^2}{8a^2}. \end{aligned}$$

Letting $\delta \rightarrow 0^+$ and $a \rightarrow \infty$ on the right hand side leads to the desired lower bound

$$(7.9) \quad \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\epsilon(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ \left. \times \gamma(B(s) - B(r)) dr ds \right\} \geq b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}.$$

□

To prove the upper bound of (7.8), it requires several steps. First, we prove

LEMMA 7.3. *Let $B_1(t)$ and $B_2(t)$ be two independent d -dimensional Brownian motions. Under $\alpha < 2$, there is a constant $C > 0$ such that*

$$(7.10) \quad \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ b t^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(s)) ds dr \right\} \leq C b^{\frac{2}{2-\alpha}}$$

for any $b > 0$.

PROOF. Perform the decomposition

$$\int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(s)) ds dr = \int_0^{ct} \int_0^{ct} + \int \int_{[0,t]^2 \setminus [0,ct]^2}$$

where the constant $0 < c < 1$ will be specified later. By Cauchy-Schwartz inequality,

$$\mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \\ \leq \left(\mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \int_0^{ct} \int_0^{ct} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \right)^{1/2} \\ \times \left(\mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \int \int_{[0,t]^2 \setminus [0,ct]^2} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \right)^{1/2}.$$

Notice that

$$\int_0^{ct} \int_0^{ct} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \\ \stackrel{d}{=} c^{\frac{4-\alpha-2\alpha_0}{2}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds.$$

Choose c such that

$$2c^{\frac{4-\alpha-2\alpha_0}{2}} = 1.$$

So we have

$$\mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \\ \leq \left(\mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \right)^{1/2} \\ \times \left(\mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \int \int_{[0,t]^2 \setminus [0,ct]^2} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \right)^{1/2}.$$

which leads to

$$(7.11) \quad \begin{aligned} & \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \\ & \leq \mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \iint_{[0,t]^2 \setminus [0,ct]^2} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\}. \end{aligned}$$

By the fact that $s+r \geq ct$ on $[0,t]^2 \setminus [0,ct]^2$,

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \iint_{[0,t]^2 \setminus [0,ct]^2} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \\ & \leq \mathbb{E}_0 \exp \left\{ \frac{2bc^{-\alpha_0}}{t} \int_0^t \int_0^t \gamma(B_1(s) + B_2(r)) dr ds \right\}. \end{aligned}$$

By (7.2), therefore,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{2b}{t^{1-\alpha_0}} \iint_{[0,t]^2 \setminus [0,ct]^2} (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \leq (2b)^{\frac{2}{2-\alpha}} \mathcal{H}.$$

In view of (7.11),

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \right\} \leq 2^{\frac{\alpha}{2-\alpha}} b^{\frac{2}{2-\alpha}} \mathcal{H}.$$

Finally, the proof follows from the identity in law:

$$\int_0^t \int_0^t (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds \stackrel{d}{=} t^{\frac{4-\alpha-\alpha_0}{2}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) dr ds.$$

□

Next, we establish a weaker version of the upper bound

LEMMA 7.4. *Under $\alpha_0 + \alpha < 2$, there is a constant $C > 0$ such that*

$$(7.12) \quad \begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ & \quad \left. \times \gamma(B(s) - B(r)) dr ds \right\} \leq C b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \end{aligned}$$

for any $b > 0$, $\theta > 0$ and $\eta > 0$.

PROOF. By the relation

$$\begin{aligned} & \int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \\ & = 2 \int \int_{[0,t]^2_{<}} \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds, \end{aligned}$$

(7.12) is equivalent to

$$(7.13) \quad \begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int \int_{[0,t]^2_{<}} \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ & \quad \left. \times \gamma(B(s) - B(r)) dr ds \right\} \leq C b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}}. \end{aligned}$$

Define the random measures $Q(\cdot)$, $Q_1(\cdot)$ and $Q_2(\cdot)$ on $(\mathbb{R}_+)_<^2$ as follows

$$Q(A) = \int \int_A \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \quad A \subset (\mathbb{R}_+)_<^2;$$

$$Q_1(A) = \int \int_A |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \quad A \subset (\mathbb{R}_+)_<^2;$$

$$Q_2(A) = \int \int_A |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \quad A \subset (\mathbb{R}_+)_<^2.$$

By the facts that

$$\left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \leq \theta^{-\alpha_0} |s-r|^{-\alpha_0}$$

and that

$$\left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \leq \eta^{-\alpha_0} |\beta(s) - \beta(r)|^{-\alpha_0}$$

we have

$$Q(A) \leq \eta^{-\alpha_0} Q_1(A) \quad \text{and} \quad Q(A) \leq \theta^{-\alpha_0} Q_2(A).$$

Consider the triangular decomposition: For a integral $N \geq 1$,

$$[0, t]_{<}^2 = \left(\bigcup_{l=0}^{2^{N+1}-1} \left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2 \right) \cup \left(\bigcup_{k=0}^{N-1} \bigcup_{l=0}^{2^k-1} A_l^k \right)$$

where


$$A_l^k = \left[\frac{2l}{2^{k+1}} t, \frac{2l+1}{2^{k+1}} t \right] \times \left[\frac{2l+1}{2^{k+1}} t, \frac{2l+2}{2^{k+1}} t \right] \quad l = 0, 1, \dots, 2^k - 1; \quad k = 0, 1, \dots, N-1.$$

See the diagram Figure 1 for the case $N = 2$. In our proof, N increases along with t in a way that will be specified later. We have

$$\begin{aligned} Q([0, t]_{<}^2) &= Q\left(\bigcup_{l=0}^{2^{N+1}-1} \left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2\right) + Q\left(\bigcup_{k=0}^{N-1} \bigcup_{l=0}^{2^k-1} A_l^k\right) \\ &\leq \eta^{-\alpha_0} Q_1\left(\bigcup_{l=0}^{2^{N+1}-1} \left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2\right) + \theta^{-\alpha_0} Q_2\left(\bigcup_{k=0}^{N-1} \bigcup_{l=0}^{2^k-1} A_l^k\right) \\ &= \eta^{-\alpha_0} \sum_{l=0}^{2^{N+1}-1} Q_1\left(\left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2\right) + \theta^{-\alpha_0} \sum_{k=0}^{N-1} Q_2\left(\bigcup_{l=0}^{2^k-1} A_l^k\right). \end{aligned}$$

By Cauchy-Schwartz inequality

$$\begin{aligned} (7.14) \quad \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} Q([0, t]_{<}^2) \right\} \\ \leq \left(\mathbb{E}_0 \exp \left\{ 2\eta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{l=0}^{2^{N+1}-1} Q_1\left(\left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2\right) \right\} \right)^{1/2} \\ \times \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{k=0}^{N-1} Q_2\left(\bigcup_{l=0}^{2^k-1} A_l^k\right) \right\} \right)^{1/2}. \end{aligned}$$

FIG 1. *triangular approximation*

Notice that the random variables

$$Q_1\left(\left[\frac{l}{2^{N+1}}t, \frac{l+1}{2^{N+1}}t\right]_{<}^2\right) \quad l = 0, 1, \dots, 2^{N+1} - 1$$

form an i.i.d. sequence with

$$\begin{aligned} Q_1\left(\left[0, \frac{t}{2^{N+1}}\right]_{<}^2\right) &= \frac{1}{2} \int_0^{\frac{t}{2^{N+1}}} \int_0^{\frac{t}{2^{N+1}}} |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \\ &\stackrel{d}{=} \frac{1}{2} \left(\frac{t}{2^{N+1}}\right)^{\frac{4-\alpha-\alpha_0}{2}} \int_0^1 \int_0^1 |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr. \end{aligned}$$

Therefore,

$$\begin{aligned} &\mathbb{E}_0 \exp \left\{ 2\eta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{l=0}^{2^{N+1}-1} Q_1\left(\left[\frac{l}{2^{N+1}}t, \frac{l+1}{2^{N+1}}t\right]_{<}^2\right) \right\} \\ &= \left(\mathbb{E}_0 \exp \left\{ \eta^{-\alpha_0} b \left(\frac{t}{2^{N+1}}\right)^{\frac{2-\alpha-\alpha_0}{2}} \frac{t^{\alpha_0}}{2^{N+1}} \int_0^1 \int_0^1 |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} \right)^{2^{N+1}}. \end{aligned}$$

Given $\epsilon > 0$, we now post our constraint

$$\eta^{-\alpha_0} \frac{t^{\alpha_0}}{2^{N+1}} \leq \epsilon \quad \text{and let} \quad N = \left\lceil \frac{\log \epsilon^{-1} (\eta^{-1} t)^{\alpha_0}}{\log 2} \right\rceil.$$

So we have

$$\begin{aligned} &\mathbb{E}_0 \exp \left\{ 2\eta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{l=0}^{2^{N+1}-1} Q_1\left(\left[\frac{l}{2^{N+1}}t, \frac{l+1}{2^{N+1}}t\right]_{<}^2\right) \right\} \\ &\leq \left(\mathbb{E}_0 \exp \left\{ \epsilon b \left(\frac{t}{2^{N+1}}\right)^{\frac{2-\alpha-\alpha_0}{2}} \int_0^1 \int_0^1 |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} \right)^{2^{N+1}}. \end{aligned}$$

Replace b by ϵb and t by $2^{-(N+1)}t$ in (7.3) and then apply it to the right hand side with the identity in law

$$\begin{aligned} & \int_0^t \int_0^t |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \\ & \stackrel{d}{=} t^{\frac{4-\alpha-\alpha_0}{2}} \int_0^1 \int_0^1 |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr. \end{aligned}$$

We have

(7.15)

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ 2\eta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{l=0}^{2^{N+1}-1} Q_1 \left(\left[\frac{l}{2^{N+1}} t, \frac{l+1}{2^{N+1}} t \right]_{<}^2 \right) \right\} \\ & \leq \lim_{t \rightarrow \infty} \frac{2^{N+1}}{t} \log \mathbb{E}_0 \exp \left\{ \epsilon b \left(\frac{t}{2^{N+1}} \right)^{\frac{2-\alpha-\alpha_0}{2}} \int_0^1 \int_0^1 |\beta(s) - \beta(r)|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} \\ & = (\epsilon b)^{\frac{2}{2-\alpha-\alpha_0}} \tilde{H} \end{aligned}$$

where we have used the fact that $2^{-(N+1)}t \rightarrow \infty$ (as $t \rightarrow \infty$) in the last equality.

Let

$$a_k = \prod_{j=2}^k \left(1 - 2^{-(1-\alpha_0)j} \right) \quad k = 2, 3, \dots, N \quad \text{and} \quad C_0 = \prod_{j=2}^{\infty} \left(1 - 2^{-(1-\alpha_0)j} \right)^{-1}$$

Since $C_0 a_N \geq 1$ and by Hölder's inequality

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\ & \leq \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_N \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\ & \leq \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_{N-1} \sum_{k=0}^{N-2} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \right)^{1-2^{-(1-\alpha_0)N}} \\ & \quad \times \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} 2^{(1-\alpha_0)N} C_0 a_N \frac{b}{t^{1-\alpha_0}} Q_2 \left(\bigcup_{l=0}^{2^N-1} A_l^{N-1} \right) \right\} \right)^{2^{-(1-\alpha_0)N}}. \end{aligned}$$

To continue, we remove the power " $1 - 2^{-(1-\alpha_0)N}$ " from the first factor. Since $a_N \leq 1$, we remove a_N from the second factor. So we have

$$\begin{aligned} (7.16) \quad & \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_N \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\ & \leq \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_{N-1} \sum_{k=0}^{N-2} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\ & \quad \times \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} 2^{(1-\alpha_0)N} C_0 \frac{b}{t^{1-\alpha_0}} Q_2 \left(\bigcup_{l=0}^{2^N-1} A_l^{N-1} \right) \right\} \right)^{2^{-(1-\alpha_0)N}}. \end{aligned}$$

Observe that

$$Q_2\left(\bigcup_{l=0}^{2^N-1} A_l^{N-1}\right) = \sum_{l=0}^{2^N-1} Q_2(A_l^{N-1})$$

and notice that the right hand side is a sum of i.i.d. Thus,

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} 2^{(1-\alpha_0)N} C_0 \frac{b}{t^{1-\alpha_0}} Q_2\left(\bigcup_{l=0}^{2^N-1} A_l^{N-1}\right) \right\} \\ &= \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} 2^{(1-\alpha_0)N} C_0 \frac{b}{t^{1-\alpha_0}} Q_2(A_0^{N-1}) \right\} \right)^{2^N}. \end{aligned}$$

Further,

$$\begin{aligned} Q_2(A_0^{N-1}) &= \int_0^{\frac{t}{2^N}} dr \int_{\frac{t}{2^N}}^{\frac{2t}{2^N}} ds |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) \\ &\stackrel{d}{=} \left(\frac{t}{2^N}\right)^{\frac{4-\alpha-2\alpha_0}{2^N}} \int_0^1 dr \int_1^2 ds |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) \\ &\stackrel{d}{=} \left(\frac{t}{2^N}\right)^{\frac{4-\alpha-2\alpha_0}{2^N}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) ds dr \end{aligned}$$

where $B_1(\cdot)$ and $B_2(\cdot)$ are two independent d -dimensional Brownian motions, and where the last identity in law follows from the variable substitution $r \mapsto 1-r$ and $s \mapsto 1+s$ so

$$\begin{aligned} & \int_0^1 dr \int_1^2 ds |s-r|^{-\alpha_0} \gamma(B(s) - B(r)) \\ &= \int_0^1 dr \int_0^1 ds |(1+s) - (1-r)|^{-\alpha_0} \gamma(B(1+s) - B(1-r)) \\ &= \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma((B(1+s) - B(1)) + (B(1) - B(1-r))) ds dr \end{aligned}$$

and $B_1(s) \equiv B(1+s) - B(1)$ ($0 \leq s \leq 1$) and $B_2(r) \equiv B(1) - B(1-r)$ ($0 \leq r \leq 1$) are two independent d -dimensional Brownian motions. In summary

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ 2^{(1-\alpha_0)N} 2\theta^{-\alpha_0} C_0 \frac{b}{t^{1-\alpha_0}} Q_2\left(\bigcup_{l=0}^{2^N-1} A_l^{N-1}\right) \right\} \\ &= \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} C_0 b \left(\frac{t}{2^N}\right)^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) ds dr \right\} \right)^{2^N}. \end{aligned}$$

Bringing it back to (7.16),

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_N \sum_{k=0}^{N-1} Q_2\left(\bigcup_{l=0}^{2^k-1} A_l^k\right) \right\} \\ &\leq \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_{N-1} \sum_{k=0}^{N-2} Q_2\left(\bigcup_{l=0}^{2^k-1} A_l^k\right) \right\} \\ &\times \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} C_0 b \left(\frac{t}{2^N}\right)^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) ds dr \right\} \right)^{2^{N\alpha_0}}. \end{aligned}$$

Repeating the same game

$$\begin{aligned}
& \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \theta^{-\alpha_0} \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\
& \leq \mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} \frac{b}{t^{1-\alpha_0}} C_0 a_N \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \\
& \leq \prod_{k=1}^N \left(\mathbb{E}_0 \exp \left\{ 2\theta^{-\alpha_0} C_0 b \left(\frac{t}{2^k} \right)^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 (s+r)^{-\alpha_0} \gamma(B_1(s) + B_2(r)) ds dr \right\} \right)^{2^{k\alpha_0}}.
\end{aligned}$$

By Lemma 7.3 with b being replaced by $2C_0 b \theta^{-\alpha_0}$ and t being replaced by $2^{-k}t$,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \theta^{-\alpha_0} \sum_{k=0}^{N-1} Q_2 \left(\bigcup_{l=0}^{2^k-1} A_l^k \right) \right\} \leq C(2C_0 b)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \sum_{k=1}^{\infty} 2^{-(1-\alpha_0)k}.$$

Together with (7.13) and (7.14),

$$\begin{aligned}
& \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} Q([0, t]_{<}^2) \right\} \\
& \leq \frac{1}{2} C(\epsilon b)^{\frac{2}{2-\alpha-\alpha_0}} + \frac{1}{2} C(2C_0 b)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \sum_{k=1}^{\infty} 2^{-(1-\alpha_0)k}.
\end{aligned}$$

Letting $\epsilon \rightarrow 0^+$ on the right hand side finally completes the proof of (7.13). $\square$

Proof of the upper bound of (7.8). To strengthen (7.12) into the upper bound requested by Theorem 7.1, we first write it as

$$\begin{aligned}
(7.17) \quad \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp & \left\{ \frac{b}{t} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\alpha_0} \right. \\
& \left. \times \gamma(B(s) - B(r)) dr ds \right\} \leq C b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}}
\end{aligned}$$

and prove

$$\begin{aligned}
(7.18) \quad \lim_{\delta \rightarrow 0^+} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp & \left\{ \frac{b}{t} \int_0^t \int_0^t \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \right. \\
& \left. \times \gamma(B(s) - B(r)) dr ds \right\} = 0
\end{aligned}$$

where

$$\tilde{\gamma}_\delta^0(u) = \Gamma(\alpha_0)^{-1} \int_{\delta^{-1}}^{\infty} e^{-\lambda|u|} \frac{d\lambda}{\lambda^{1-\alpha_0}}.$$

Indeed, given $\epsilon > 0$, break $[0, t]^2$ into two parts: On the first part

$$\left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right| \geq \epsilon$$

where

$$\tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \leq \tilde{\gamma}_\delta^0(\epsilon).$$

On the second part

$$\left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right| < \epsilon$$

where

$$\begin{aligned} \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) &\leq \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\alpha_0} \\ &\leq \epsilon^{\tilde{\alpha}_0 - \alpha_0} \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\tilde{\alpha}_0} \end{aligned}$$

and where $\alpha_0 < \tilde{\alpha}_0 < 1$ satisfies $\tilde{\alpha}_0 + \alpha < 2$. Thus,

$$\begin{aligned} &\int_0^t \int_0^t \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \gamma(B(s) - B(r)) dr ds \\ &\leq \tilde{\gamma}_\delta^0(\epsilon) \int_0^t \int_0^t \gamma(B(s) - B(r)) dr ds \\ &\quad + \epsilon^{\tilde{\alpha}_0 - \alpha_0} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\tilde{\alpha}_0} \gamma(B(s) - B(r)) dr ds. \end{aligned}$$

By Cauchy-Schwartz's inequality

$$\begin{aligned} &\mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \\ &\leq \left(\mathbb{E}_0 \exp \left\{ \frac{2b}{t} \tilde{\gamma}_\delta^0(\epsilon) \int_0^t \int_0^t \gamma(B(s) - B(r)) dr ds \right\} \right)^{1/2} \\ &\quad \times \left(\mathbb{E}_0 \exp \left\{ \frac{2b}{t} \epsilon^{\tilde{\alpha}_0 - \alpha_0} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\tilde{\alpha}_0} \gamma(B(s) - B(r)) dr ds \right\} \right)^{1/2}. \end{aligned}$$

By (7.2) and (7.17) (with α_0 being replaced by $\tilde{\alpha}_0$),

$$\begin{aligned} &\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \\ &\leq \frac{1}{2} \left\{ C(2b)^{\frac{2}{2-\alpha}} (\tilde{\gamma}_\delta^0(\epsilon))^{\frac{2}{2-\alpha}} + C(2b)^{\frac{2}{2-\alpha}} \epsilon^{\frac{2(\alpha_0 - \alpha_0)}{2-\alpha}} \right\}. \end{aligned}$$

The claim (7.18) follows from the fact that $\tilde{\gamma}_\delta^0(\epsilon) \rightarrow 0$ ($\delta \rightarrow 0^+$) for any $\epsilon > 0$.

We finally come to the proof for the upper bound of (7.8). By the relation $|\cdot|^{-\alpha_0} = \gamma_\delta^0(\cdot) + \tilde{\gamma}_\delta^0(\cdot)$ and by Hölder's inequality

$$\begin{aligned} &\mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ &\leq \left(\mathbb{E}_0 \exp \left\{ \frac{bp}{t} \int_0^t \int_0^t \gamma_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \right)^{1/p} \\ &\quad \times \left(\mathbb{E}_0 \exp \left\{ \frac{bq}{t} \int_0^t \int_0^t \tilde{\gamma}_\delta^0 \left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right) \gamma(B(s) - B(r)) dr ds \right\} \right)^{1/q}. \end{aligned}$$

Using the bound

$$\gamma_\delta^0\left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t}\right) \leq \gamma_\delta^0\left(\frac{\theta(s-r)}{t}\right) = \theta^{-\alpha_0} \gamma_{\theta^{-1}\delta}\left(\frac{s-r}{t}\right)$$

and (7.7) (with b being replaced by $bp\theta^{-\alpha_0}$ and δ by $\theta^{-1}\delta$),

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{bp}{t} \int_0^t \int_0^t \gamma_\delta^0\left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t}\right) \gamma(B(s) - B(r)) dr ds \right\} \\ & \leq (pb)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}_{\theta^{-1}\delta} \leq (pb)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}. \end{aligned}$$

Thus,

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ & \leq \frac{1}{p} (pb)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E} \\ & + \frac{1}{q} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{bq}{t} \int_0^t \int_0^t \tilde{\gamma}_\delta^0\left(\frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t}\right) \gamma(B(s) - B(r)) dr ds \right\}. \end{aligned}$$

Letting $\delta \rightarrow 0^+$ on the right hand side, by (7.18) (with b being replaced by qb),

$$\begin{aligned} & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t} \int_0^t \int_0^t \left| \frac{\theta(s-r) + i\eta(\beta(s) - \beta(r))}{t} \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ & \leq \frac{1}{p} (pb)^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}. \end{aligned}$$

Letting $p \rightarrow 1^+$ on the right hand side leads to the demanded upper bound:

$$\begin{aligned} (7.19) \quad & \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ \frac{b}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ & \quad \left. \times \gamma(B(s) - B(r)) dr ds \right\} \leq b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}. \end{aligned}$$

□

To be applied to the proof of Theorem 1.2, the large deviations discussed in this section have to be transformed from the forms of exponential moments into the forms of polynomial moments. We end this section with the following lemma.

LEMMA 7.5. *Recall that $\mathcal{E}$, $\mathcal{E}_*$ and $\gamma_*^0(\cdot)$ are introduced in Theorem 1.2. We have*

$$\begin{aligned} (7.20) \quad & \lim_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{-\alpha/2} \mathbb{E}_0 \left[\int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \right]^n \\ & = \log \left(\frac{2\mathcal{E}_*}{2-\alpha} \right)^{\frac{2-\alpha}{2}}. \end{aligned}$$

In addition,

$$\begin{aligned} (7.21) \quad & \lim_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{-\alpha/2} \mathbb{E}_0 \left[\int_0^1 \int_0^1 \left| \theta(s-r) + i \frac{\eta}{\sqrt{n}} (\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ & \quad \left. \times \gamma(B(s) - B(r)) ds dr \right]^n \\ & = \log \left(\frac{2\mathcal{E}}{2-\alpha} \right)^{\frac{2-\alpha}{2}} - \alpha_0 \log \theta. \end{aligned}$$

for any $\theta > 0$ and $\eta > 0$,

PROOF. By the bound

$$\begin{aligned}\gamma_*^0(u) &\leq \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+} \exp\left\{-\frac{\lambda^2}{2(4-\alpha-2\alpha_0)}|u|\right\} \frac{d\lambda}{\lambda^{1-\alpha_0}} \\ &= \frac{1}{2} \Gamma(\alpha_0)^{-1} \Gamma\left(\frac{\alpha_0}{2}\right) \left(\frac{|u|}{2(4-\alpha-2\alpha_0)}\right)^{-\frac{\alpha_0}{2}}\end{aligned}$$

the condition (7.5) holds for $\tilde{\gamma}^0(\cdot) = \gamma_*^0(\cdot)$ under $\alpha_0 + \alpha < 2$. By scaling,

$$\int_0^t \int_0^t \gamma_*^0\left(\frac{s-r}{t}\right) \gamma(B(s) - B(r)) ds dr \stackrel{d}{=} t^{\frac{4-\alpha}{2}} \int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr.$$

Therefore, (7.6) can be written as

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ b t^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \right\} = b^{\frac{2}{2-\alpha}} \mathcal{E}_* \quad b > 0.$$

By Gärtner-Ellis theorem on $\mathbb{R}^+$ (Theorem 1.2.4, p. 11, [5]), the process

$$t^{-\alpha/2} \int_0^t \int_0^t \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \quad t > 0$$

obeys the large deviation on $\mathbb{R}_+$ with deviation scale t and the rate function

$$I(\lambda) = \sup_{b>0} \left\{ \lambda b - b^{\frac{2}{2-\alpha}} \mathcal{E}_* \right\} = \frac{\alpha}{2} \left(\frac{2-\alpha}{2\mathcal{E}_*} \right)^{\frac{2-\alpha}{\alpha}} \lambda^{\frac{2}{\alpha}} \quad \lambda > 0.$$

By Varadhan's integral lemma (Theorem 4.3, p.137, [17])

$$\begin{aligned}&\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ t \log \left(t^{-\alpha/2} \int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \right) \right\} \\ &= \sup_{\lambda > 0} \left\{ \log \lambda - \frac{\alpha}{2} \left(\frac{2-\alpha}{2\mathcal{E}_*} \right)^{\frac{2-\alpha}{\alpha}} \lambda^{2/\alpha} \right\} = -\frac{\alpha}{2} + \log \left(\frac{2\mathcal{E}_*}{2-\alpha} \right)^{\frac{2-\alpha}{2}}.\end{aligned}$$

Take $t = n$:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log n^{-\frac{\alpha n}{2}} \mathbb{E}_0 \left[\int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \right]^n = -\frac{\alpha}{2} + \log \left(\frac{2\mathcal{E}_*}{2-\alpha} \right)^{\frac{2-\alpha}{2}}.$$

Applying Stirling formula to the above gives (7.20).

Notice that

$$\begin{aligned}&\int_0^t \int_0^t \left| \theta(s-r) + i\eta(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \\ &\stackrel{d}{=} t^{\frac{4-\alpha-2\alpha_0}{2}} \int_0^1 \int_0^1 \left| \theta(s-r) + i\frac{\eta}{\sqrt{t}}(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds.\end{aligned}$$

Therefore, (7.8) can be written as

$$\begin{aligned}&\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}_0 \exp \left\{ b t^{\frac{2-\alpha}{2}} \int_0^1 \int_0^1 \left| \theta(s-r) + i\frac{\eta}{\sqrt{t}}(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) dr ds \right\} \\ &= b^{\frac{2}{2-\alpha}} \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E}.\end{aligned}$$

The remaining of the proof for (7.21) follows a completely parallel argument. $\square$

8. Proof of Theorem 1.2. The central piece of the proof is to establish

$$\begin{aligned}
 (8.1) \quad & \log \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}_*}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right) \\
 & \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{3 - \alpha} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\
 & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{3 - \alpha} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\
 & \leq \log \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right).
 \end{aligned}$$

Indeed, by (2.20) $\mathbb{E} S_{2n-1}(g_{2n-1}(\cdot, t, 0)) = 0$. From (2.22), (1.8) and (2.6) one can derive that

$$(8.2) \quad \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) = t^{(4 - \alpha - \alpha_0)n} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \quad t > 0.$$

By the Stratonovich expansion (2.13), therefore,

$$\mathbb{E} u(t, x) = \mathbb{E} u(t, 0) = \sum_{n=0}^{\infty} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) = \sum_{n=0}^{\infty} t^{(4 - \alpha - \alpha_0)n} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0))$$

where the first equality comes from the stationarity in x . By (8.1), therefore, the upper bound of Theorem 1.2 follows from the following computation:

$$\begin{aligned}
 & \limsup_{t \rightarrow \infty} t^{-\frac{4 - \alpha - \alpha_0}{3 - \alpha}} \log \mathbb{E} u(t, x) = \limsup_{t \rightarrow \infty} t^{-\frac{4 - \alpha - \alpha_0}{3 - \alpha}} \log \sum_{n=0}^{\infty} t^{(4 - \alpha - \alpha_0)n} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\
 & = \lim_{t \rightarrow \infty} t^{-\frac{4 - \alpha - \alpha_0}{3 - \alpha}} \log \sum_{n=0}^{\infty} \frac{t^{(4 - \alpha - \alpha_0)n}}{(n!)^{3 - \alpha}} \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right)^n \\
 & = (3 - \alpha) \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right)
 \end{aligned}$$

where the last step follows from the elementary identity on asymptotics of Mittag-Leffler function (Lemma A.3, [4]):

$$\lim_{b \rightarrow \infty} b^{-1/\gamma} \log \sum_{n=0}^{\infty} \frac{\theta^n b^n}{(n!)^\gamma} = \gamma \theta^{1/\gamma} \quad \theta, \gamma > 0$$

with $b = t^{4 - \alpha - \alpha_0}$, $\gamma = 3 - \alpha$ and

$$\theta = \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right)$$

Similarly, from (8.1) we get the lower bound

$$\liminf_{t \rightarrow \infty} t^{-\frac{4 - \alpha - \alpha_0}{3 - \alpha}} \log \mathbb{E} u(t, x) \geq (3 - \alpha) \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4 - \alpha - 2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4 - \alpha - \alpha_0}} \left(\frac{2\mathcal{E}_*}{2 - \alpha} \right)^{\frac{2 - \alpha}{2}} \right)$$

In the following subsections, we shall establish the upper and lower bounds, separately, for (8.1).

8.1. *Upper bound for (8.1).* In view of (6.1) and (7.1),

$$(8.3) \quad \int_0^\infty e^{-\theta t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt \\ \leq \frac{\theta}{2} \left(\frac{1}{2}\right)^{3n} \frac{1}{n!} \int_0^\infty dt \exp\left\{-\frac{\theta^2}{2}t\right\} \\ \times \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n.$$

Let $\theta > 0$ be fixed but arbitrary (for a while). Given $\eta > 0$, write

$$(8.4) \quad \int_0^\infty dt \exp\left\{-\frac{\theta^2}{2}t\right\} \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \right. \\ \left. \times \gamma(B(s) - B(r)) ds dr \right]^n \\ = \left\{ \int_0^{\eta^{-2}n} + \int_{\eta^{-2}n}^\infty \right\} dt \exp\left\{-\frac{\theta^2}{2}t\right\} \\ \times \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n.$$

For the first term on the right hand side, we use the scaling identity

$$\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \\ \stackrel{d}{=} t^{\frac{4-\alpha-2\alpha_0}{2}} \int_0^1 \int_0^1 \left| \theta(s-r) + i \frac{1}{\sqrt{t}} (\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr.$$

So we have

$$\int_0^{\eta^{-2}n} dt \exp\left\{-\frac{\theta^2}{2}t\right\} \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ = \int_0^{\epsilon^{-2}n} dt \exp\left\{-\frac{\theta^2}{2}t\right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} \\ \times \mathbb{E}_0 \left[\int_0^1 \int_0^1 \left| \theta(s-r) + i \frac{1}{\sqrt{t}} (\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ \leq \mathbb{E}_0 \left[\int_0^1 \int_0^1 \left| \theta(s-r) + i \frac{\eta}{\sqrt{n}} (\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ \times \int_0^\infty \exp\left\{-\frac{\theta^2}{2}t\right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} dt \\ = \mathbb{E}_0 \left[\int_0^1 \int_0^1 \left| \theta(s-r) + i \frac{\eta}{\sqrt{n}} (\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ \times \left(\frac{2}{\theta^2}\right)^{1+\frac{4-\alpha-2\alpha_0}{2}n} \Gamma\left(1 + \frac{4-\alpha-2\alpha_0}{2}n\right) \\ = (1+o(1))^n (n!)^{2-\alpha_0} (4-\alpha-2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}n} \theta^{-(4-\alpha-\alpha_0)n} \left(\frac{2\mathcal{E}}{2-\alpha}\right)^{\frac{2-\alpha}{2}n} \quad (n \rightarrow \infty)$$

where the last step follows from (7.21) and Stirling formula.

As for the second term, we use the bound of Taylor expansion

$$\begin{aligned} & \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ & \leq n! t^{(1-\alpha_0)n} \mathbb{E}_0 \exp \left\{ \frac{1}{t^{1-\alpha_0}} \int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right\} \\ & \leq (1 + o(1))^n n! t^{(1-\alpha_0)n} \exp \left\{ \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E} t \right\} \quad (n \rightarrow \infty) \end{aligned}$$

for large t , where the last step follows from Theorem 7.1 with $b = \eta = 1$. Take $\theta > 0$ sufficiently large so that

$$c(\theta) \equiv \frac{\theta^2}{2} - \theta^{-\frac{2\alpha_0}{2-\alpha}} \mathcal{E} > 0.$$

We have

$$\begin{aligned} & \int_{\eta^{-2n}}^{\infty} dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \mathbb{E}_0 \left[\int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ & \leq (1 + o(1))^n n! \int_{\eta^{-2n}}^{\infty} t^{(1-\alpha_0)n} \exp \{-c(\theta)t\} dt \quad (n \rightarrow \infty). \end{aligned}$$

We now claim that

$$(8.5) \quad \lim_{\eta \rightarrow 0^+} \limsup_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{-(1-\alpha_0)} \int_{\eta^{-2n}}^{\infty} t^{(1-\alpha_0)n} \exp \{-c(\theta)t\} dt = -\infty.$$

Indeed, consider an i.i.d. sequence $X_1, \dots, X_n$ with common distribution $\Gamma(1 - \alpha_0, c(\theta))$, i.e., they have the common density

$$f(x) = \Gamma(1 - \alpha_0)^{-1} c(\theta)^{(1-\alpha_0)} x^{-\alpha_0} \exp \left\{ -c(\theta)x \right\} \quad x > 0$$

and $X_0 \sim \exp(c(\theta))$ independent of $\{X_1, \dots, X_n\}$. We have that

$$X_0 + X_1 + \dots + X_n \sim \Gamma(1 + n(1 - \alpha_0), c(\theta)).$$

Therefore

$$\begin{aligned} & \Gamma(1 + n(1 - \alpha_0))^{-1} c(\theta)^{1+(1-\alpha_0)n} \int_{\eta^{-2n}}^{\infty} t^{(1-\alpha_0)n} \exp \{-c(\theta)t\} dt \\ & = \mathbb{P} \left\{ \frac{X_0 + X_1 + \dots + X_n}{n} \geq \eta^{-2} \right\}. \end{aligned}$$

Therefore, (8.5) follows from Cramér large deviation (Theorem 2.2.3, p.27, [17]) which particularly leads to

$$\lim_{\eta \rightarrow 0^+} \limsup_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{P} \left\{ \frac{X_0 + X_1 + \dots + X_n}{n} \geq \eta^{-2} \right\} = -\infty.$$

In summary,

$$\begin{aligned} & \int_{\eta^{-2n}}^{\infty} dt \exp \left\{ -\frac{\theta^2}{2} t \right\} \int_0^t \int_0^t \left| \theta(s-r) + i(\beta(s) - \beta(r)) \right|^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \Big]^n \\ & \leq (1 + o(1))^n (n!)^{2-\alpha_0} \exp \{-L_\eta n\} \end{aligned}$$

where $L_\eta > 0$ can be sufficiently large if η is sufficiently small.

So we reach the point that in the decomposition (8.4), the bound of the first term dominates the bound of the second term as $\eta > 0$ is small. Consequently, by (8.3) we have

$$\begin{aligned} & \int_0^\infty e^{-\theta t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt \\ & \leq (1 + o(1))^n \left(\frac{1}{2}\right)^{3n} (n!)^{1-\alpha_0} (4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}n} \theta^{-(4-\alpha-\alpha_0)n} \left(\frac{2\mathcal{E}}{2-\alpha}\right)^{\frac{2-\alpha}{2}n}. \end{aligned}$$

On the other hand, by (8.2)

$$\begin{aligned} & \int_0^\infty e^{-\theta t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt = \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \int_0^\infty e^{-\theta t} t^{(4-\alpha-\alpha_0)n} dt \\ & = \theta^{-(1+(4-\alpha-\alpha_0)n)} \Gamma(1 + (4 - \alpha - \alpha_0)n) \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\ & = (1 + o(1))^n \theta^{-(4-\alpha-\alpha_0)n} (4 - \alpha - \alpha_0)^{(4-\alpha-\alpha_0)n} (n!)^{4-\alpha-\alpha_0} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)). \end{aligned}$$

So we have the upper bound of (8.1):

$$\begin{aligned} (8.6) \quad & \limsup_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{3-\alpha} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\ & \leq \log \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4-\alpha-\alpha_0}} \left(\frac{2\mathcal{E}}{2-\alpha}\right)^{\frac{2-\alpha}{2}} \right). \end{aligned}$$

□

8.2. *Lower bound for (8.1).* Take $\theta = 1$. Related to the representation (6.1) and by (6.3)

$$\begin{aligned} & \int_0^t \int_0^t \left(((s-r) + i(\beta(s) - \beta(r)) \operatorname{sgn}(s-r)) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \\ & = \Gamma(\alpha_0)^{-1} \int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^t \int_0^t \exp \left\{ -|s-r| + i(\beta(s) - \beta(r)) \operatorname{sgn}(s-r) \right\} \\ & \quad \times \gamma(B(s) - B(r)) ds dr. \end{aligned}$$

Hence

$$\begin{aligned} & \mathbb{E}_0 \left[\int_0^t \int_0^t \left(((s-r) + i(\beta(s) - \beta(r)) \operatorname{sgn}(s-r)) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\ & = \int_{\mathbb{R}_+^n} \left(\prod_{l=1}^n \frac{d\lambda_l}{\lambda_l^{1-\alpha_0}} \right) \int_{[0,t]^{2n}} dr_1 ds_1 \cdots dr_n ds_n \exp \left\{ -\theta \sum_{l=1}^n \lambda_l |s_l - r_l| \right\} \\ & \quad \times \mathbb{E}_0 \exp \left\{ i \sum_{l=1}^n \lambda_l (\beta(s_l) - \beta(r_l)) \operatorname{sgn}(s_l - r_l) \right\} \mathbb{E}_0 \prod_{l=1}^n \gamma(B(s_l) - B(r_l)). \end{aligned}$$

Notice that

$$\begin{aligned} & \mathbb{E}_0 \exp \left\{ i \sum_{l=1}^n \lambda_l (\beta(s_l) - \beta(r_l)) \operatorname{sgn}(s_l - r_l) \right\} \\ & = \exp \left\{ -\frac{1}{2} \operatorname{Var} \left(\sum_{l=1}^n \lambda_l (\beta(s_l) - \beta(r_l)) \operatorname{sgn}(s_l - r_l) \right) \right\} \\ & \geq \exp \left\{ -\frac{n}{2} \sum_{l=1}^n \lambda_l^2 \operatorname{Var}(\beta(s_l) - \beta(r_l)) \right\} = \exp \left\{ -\frac{n}{2} \sum_{l=1}^n \lambda_l^2 |s_l - r_l| \right\}. \end{aligned}$$

In summary,

(8.7)

$$\begin{aligned}
& \mathbb{E}_0 \left[\int_0^t \int_0^t \left(((s-r) + i(\beta(s) - \beta(r)) \operatorname{sgn}(s-r)) \right)^{-\alpha_0} \gamma(B(s) - B(r)) ds dr \right]^n \\
& \geq \Gamma(\alpha_0)^{-n} \int_{\mathbb{R}_+^n} \left(\prod_{l=1}^n \frac{d\lambda_l}{\lambda_l^{1-\alpha_0}} \right) \int_{[0,t]^{2n}} dr_1 ds_1 \cdots dr_n ds_n \\
& \quad \times \exp \left\{ - \sum_{l=1}^n \left(\lambda_l + \frac{n\lambda_l^2}{2} \right) |s_l - r_l| \right\} \mathbb{E}_0 \prod_{l=1}^n \gamma(B(s_l) - B(r_l)) \\
& = \Gamma(\alpha_0)^{-n} \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^t \int_0^t \exp \left\{ - \left(\lambda + \frac{n\lambda^2}{2} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n
\end{aligned}$$

By scaling

$$\begin{aligned}
& \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^t \int_0^t \exp \left\{ - \left(\lambda + \frac{n\lambda^2}{2} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n \\
& = t^{\frac{4-\alpha-2\alpha_0}{2}n} \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^1 \int_0^1 \exp \left\{ - \left(\lambda + \frac{n\lambda^2}{2t} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n.
\end{aligned}$$

By Theorem 6.1 with $\theta = 1$, therefore,

$$\begin{aligned}
& \int_0^\infty e^{-t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt \\
& \geq \Gamma(\alpha_0)^{-n} \frac{1}{2} \left(\frac{1}{2} \right)^{3n} \frac{1}{n!} \int_0^\infty dt \exp \left\{ - \frac{t}{2} \right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} \\
& \quad \times \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^1 \int_0^1 \exp \left\{ - \left(\lambda + \frac{n\lambda^2}{2t} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n \\
& \geq \Gamma(\alpha_0)^{-n} \frac{1}{2} \left(\frac{1}{2} \right)^{3n} \frac{1}{n!} \left(\int_{nc}^\infty \exp \left\{ - \frac{t}{2} \right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} dt \right) \\
& \quad \times \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^1 \int_0^1 \exp \left\{ - \left(\lambda + \frac{\lambda^2}{2c} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n
\end{aligned}$$

where $c = 4 - \alpha - \alpha_0$. Notice that

$$\begin{aligned}
& \Gamma(\alpha_0)^{-n} \mathbb{E}_0 \left[\int_{\mathbb{R}_+} \frac{d\lambda}{\lambda^{1-\alpha_0}} \int_0^1 \int_0^1 \exp \left\{ - \left(\lambda + \frac{\lambda^2}{2c} \right) |s-r| \right\} \gamma(B(s) - B(r)) ds dr \right]^n \\
& = \mathbb{E}_0 \left[\int_0^1 \int_0^1 \gamma_*^0(s-r) \gamma(B(s) - B(r)) ds dr \right]^n.
\end{aligned}$$

By constructing relevant independent Gamma-distributed random variables (as we did in the proof of (8.5)) and by the central limit theorem, one can show that

$$\begin{aligned}
& \int_{nc}^\infty \exp \left\{ - \frac{t}{2} \right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} dt \sim \frac{1}{2} \int_0^\infty \exp \left\{ - \frac{t}{2} \right\} t^{\frac{4-\alpha-2\alpha_0}{2}n} dt \\
& = 2^{\frac{4-\alpha-2\alpha_0}{2}n} \Gamma \left(1 + \frac{4-\alpha-2\alpha_0}{2}n \right) \quad (n \rightarrow \infty).
\end{aligned}$$

By (7.20) and Stirling formula, therefore,

$$\begin{aligned} & \int_0^\infty e^{-t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt \\ & \geq (1 + o(1))^n \left(\frac{1}{2}\right)^{3n} (n!)^{1-\alpha_0} (4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}n} \left(\frac{2\mathcal{E}_*}{2-\alpha}\right)^{\frac{2-\alpha}{2}n} \end{aligned}$$

as $n \rightarrow \infty$.

On the other hand, by (8.2)

$$\begin{aligned} & \int_0^\infty e^{-t} \mathbb{E} S_{2n}(g_{2n}(\cdot, t, 0)) dt = \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \int_0^\infty e^{-t} t^{(4-\alpha-\alpha_0)n} dt \\ & = \Gamma\left(1 + (4 - \alpha - \alpha_0)n\right) \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\ & = (1 + o(1))^n (n!)^{4-\alpha-\alpha_0} (4 - \alpha - \alpha_0)^{(4-\alpha-\alpha_0)n} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)). \end{aligned}$$

Combining them together, we have completed the proof of the lower bound of (8.1).

$$\begin{aligned} (8.8) \quad & \liminf_{n \rightarrow \infty} \frac{1}{n} \log(n!)^{3-\alpha} \mathbb{E} S_{2n}(g_{2n}(\cdot, 1, 0)) \\ & \geq \log \left(\frac{(4 - \alpha - 2\alpha_0)^{\frac{4-\alpha-2\alpha_0}{2}}}{2^3(4 - \alpha - \alpha_0)^{4-\alpha-\alpha_0}} \left(\frac{2\mathcal{E}_*}{2-\alpha}\right)^{\frac{2-\alpha}{2}} \right). \end{aligned}$$

□

REMARK 8.1. In view of our proof for the lower bound, it seems that (8.8) is not sharp to the constant as we are not able to undo the impact of " $\beta(s) - \beta(r)$ " in the representation (6.1). The challenge comes from the fact that the Brownian intersection local time appearing on the right hand side of (6.1) takes complex values. As a consequence, the usual treatments for the lower bound, such as Jansen's inequality, can not be applied here.

APPENDIX

Let $\theta > 0$. Corresponding to the norms $\|\cdot\|_{j,k}^{(1)}$ and $\|\cdot\|_{j,k}^{(2)}$ introduced in (3.4) and (3.5), the following lemma is concerned with the bound given in Theorem 3.1 with $G_l(t, x) = e^{-\theta t} G(t, x)$.

LEMMA A.1. Assume (1.5).

(i) For any $\theta > 0$,

$$\begin{aligned} (A.1) \quad & \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \int_0^\infty \int_{\mathbb{R}^d} e^{-(\theta+\lambda)t+i\xi \cdot x} G(t, x) dx dt \right| \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \\ & = \int_{\mathbb{R}_+ \times \mathbb{R}^d} \frac{1}{(\theta + \lambda)^2 + |\xi|^2} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) < \infty. \end{aligned}$$

(ii) For any $\theta > 0$,

$$\begin{aligned} (A.2) \quad & \int_0^\infty \int_0^\infty ds dt e^{-\theta(t+s)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |s - t|^{-\alpha_0} \gamma(x - y) G(t, x) G(s, y) dx dy \\ & = \frac{1}{2} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \frac{1}{(\theta + \lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\theta} \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \frac{\theta + \lambda}{(\theta + \lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2} \\
& < \infty.
\end{aligned}$$

Further, there is a constant $C > 0$ such that

$$(A.3) \quad \int_0^\infty \int_0^\infty ds dt e^{-\theta(t+s)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |s-t|^{-\alpha_0} \gamma(x-y) G(t,x) G(s,y) dx dy \leq C \theta^{-2}$$

for large θ .

PROOF. The identity in (A.1) follows from

$$\int_0^\infty \int_{\mathbb{R}^d} e^{-(\theta+\lambda)t + i\xi \cdot x} G(t,x) dx dt = \int_0^\infty e^{-(\theta+\lambda)t} \frac{\sin(t|\xi|)}{|\xi|} dt = \frac{1}{(\theta + \lambda)^2 + |\xi|^2}$$

where the first equality follows from (2.3) and the second from integration by parts. To show the finiteness,

$$\int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{1}{(\theta + \lambda)^2 + |\xi|^2} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \leq \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{1}{\lambda^2 + (\theta^2 + |\xi|^2)} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi)$$

and by variable substitution

$$\int_{\mathbb{R}^+} \frac{1}{\lambda^2 + (\theta^2 + |\xi|^2)} \frac{d\lambda}{\lambda^{1-\alpha_0}} = \left(\frac{1}{\theta^2 + |\xi|^2} \right)^{\frac{2-\alpha_0}{2}} \int_{\mathbb{R}^+} \frac{1}{1 + \lambda^2} \frac{d\lambda}{\lambda^{1-\alpha_0}}.$$

By (1.5), therefore,

$$\int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{1}{(\theta + \lambda)^2 + |\xi|^2} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \leq C \int_{\mathbb{R}^d} \left(\frac{1}{\theta^2 + |\xi|^2} \right)^{\frac{2-\alpha_0}{2}} \mu(d\xi) < \infty.$$

We now prove (A.2). By (1.5),

$$\begin{aligned}
& \int_0^\infty \int_0^\infty ds dt e^{-\theta(t+s)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |s-t|^{-\alpha_0} \gamma(x-y) G(t,x) G(s,y) dx dy \\
& = 2 \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \int \int_{\{s \leq t\}} ds dt e^{-\theta(s+t)} e^{-\lambda(t-s)} \int_{\mathbb{R}^d \times \mathbb{R}^d} e^{i\xi \cdot (x-y)} G(t,x) G(s,y) dx dy \\
& = 2 \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \int \int_{\{s \leq t\}} ds dt e^{-\lambda(t-s)} e^{-\theta(s+t)} \frac{\sin(t|\xi|) \sin(s|\xi|)}{|\xi|^2} \\
& = 2 \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \frac{\mu(d\xi)}{|\xi|^2} \int_0^\infty ds e^{-2\theta s} \sin(s|\xi|) \int_s^\infty e^{-(\theta+\lambda)(t-s)} \sin(t|\xi|) dt.
\end{aligned}$$

By the relation

$$\sin(t|\xi|) = \sin((t-s)|\xi|) \cos(s|\xi|) + \cos((t-s)|\xi|) \sin(s|\xi|)$$

and by integration by parts

$$\begin{aligned}
& \int_s^\infty e^{-(\theta+\lambda)(t-s)} \sin(t|\xi|) dt \\
& = \cos(s|\xi|) \int_0^\infty e^{-(\theta+\lambda)t} \sin(t|\xi|) dt + \sin(s|\xi|) \int_0^\infty e^{-(\theta+\lambda)t} \cos(t|\xi|) dt \\
& = \cos(s|\xi|) \frac{|\xi|}{(\theta + \lambda)^2 + |\xi|^2} + \sin(s|\xi|) \frac{\theta + \lambda}{(\theta + \lambda)^2 + |\xi|^2}.
\end{aligned}$$

Thus,

$$\begin{aligned} & \int_0^\infty \int_0^\infty ds dt e^{-\theta(t+s)} \int_{\mathbb{R}^d \times \mathbb{R}^d} |s-t|^{-\alpha_0} \gamma(x-y) G(t,x) G(s,y) dx dy \\ &= 2 \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \frac{\mu(d\xi)}{|\xi|^2} \frac{|\xi|}{(\theta+\lambda)^2 + |\xi|^2} \int_0^\infty e^{-2\theta s} \sin(s|\xi|) \cos(s|\xi|) ds \\ &+ 2 \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \frac{\mu(d\xi)}{|\xi|^2} \frac{\theta+\lambda}{(\theta^2+\lambda)^2 + |\xi|^2} \int_0^\infty e^{-2\theta s} \sin^2(s|\xi|) ds. \end{aligned}$$

Using integration by parts again

$$\int_0^\infty e^{-2\theta s} \sin(s|\xi|) \cos(s|\xi|) ds = \frac{1}{2} \int_0^\infty e^{-2\theta s} \sin(2s|\xi|) ds = \frac{1}{2} \frac{2|\xi|}{4\theta^2 + 4|\xi|^2} = \frac{1}{4} \frac{|\xi|}{\theta^2 + |\xi|^2}.$$

A similar treatment also leads to

$$\int_0^\infty e^{-2\theta s} \sin^2(s|\xi|) ds = \frac{|\xi|}{2\theta} \int_0^\infty e^{-2\theta s} \sin(2s|\xi|) ds = \frac{1}{4\theta} \frac{|\xi|^2}{\theta^2 + |\xi|^2}.$$

Bringing them together leads to the identity leads to the identity in (A.2).

Establishing the finiteness in (A.2) is an easy job and can be seen from the following estimate for (A.3). To show (A.3), all we need is to bound the two terms on the right hand side of (A.2) separately. We first work on the second term.

By variable substitution

$$\begin{aligned} & \int_0^\infty \frac{d\lambda}{\lambda^{1-\alpha_0}} \frac{\theta+\lambda}{(\theta+\lambda)^2 + |\xi|^2} = \int_\theta^\infty \frac{d\lambda}{(\lambda-\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + |\xi|^2} \\ &= \frac{1}{|\xi|^{1-\alpha_0}} \int_{|\xi|^{-1}\theta}^\infty \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1}. \end{aligned}$$

Consider the decomposition

$$\int_{|\xi|^{-1}\theta}^\infty \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1} \leq \left\{ \int_{|\xi|^{-1}\theta}^{2|\xi|^{-1}\theta} + \int_{2|\xi|^{-1}\theta}^\infty \right\} \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1}.$$

For the first term

$$\begin{aligned} & \int_{|\xi|^{-1}\theta}^{2|\xi|^{-1}\theta} \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1} \leq \left(\frac{|\xi|}{\theta} \right) \int_{|\xi|^{-1}\theta}^{2|\xi|^{-1}\theta} \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \\ &= \left(\frac{|\xi|}{\theta} \right) \int_0^{|\xi|^{-1}\theta} \frac{d\lambda}{\lambda^{1-\alpha_0}} = \frac{1}{\alpha_0} \left(\frac{|\xi|}{\theta} \right)^{1-\alpha_0}. \end{aligned}$$

As for the second term

$$\int_{2|\xi|^{-1}\theta}^\infty \frac{d\lambda}{(\lambda - |\xi|^{-1}\theta)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1} \leq \int_{2|\xi|^{-1}\theta}^\infty \frac{d\lambda}{(\lambda/2)^{1-\alpha_0}} \frac{\lambda}{\lambda^2 + 1} \leq C \left(\frac{|\xi|}{\theta} \right)^{1-\alpha_0}.$$

In summary, we have the bound (the constant C can be different from place to place in our argument)

$$\int_0^\infty \frac{d\lambda}{\lambda^{1-\alpha_0}} \frac{\theta+\lambda}{(\theta+\lambda)^2 + |\xi|^2} \leq C \theta^{-(1-\alpha_0)}.$$

By Fubini's theorem, therefore,

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \frac{\theta+\lambda}{(\theta+\lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2} \leq C \theta^{-(1-\alpha_0)} \int_{\mathbb{R}^d} \frac{1}{\theta^2 + |\xi|^2} \mu(d\xi) \\ &\leq C \theta^{-(3-\alpha_0)} \mu(\{|\xi| \leq 1\}) + C \theta^{-(1-\alpha_0)} \int_{\{|\xi| \geq 1\}} \frac{1}{\theta^2 + |\xi|^2} \mu(d\xi). \end{aligned}$$

By Minkowski inequality with $p = 2/\alpha_0$ and $q = 2(2 - \alpha_0)^{-1}$,

$$\theta^2 + |\xi|^2 \geq C^{-1} \theta^{\alpha_0} |\xi|^{2-\alpha_0}.$$

So we have

$$\theta^{-(1-\alpha_0)} \int_{\{|\xi| \geq 1\}} \frac{1}{\theta^2 + |\xi|^2} \mu(d\xi) \leq C \theta^{-1} \int_{\{|\xi| \geq 1\}} |\xi|^{-(2-\alpha_0)} \mu(d\xi).$$

The integral on the right hand side is finite under (1.5). Therefore, we have established the expected bound

$$\frac{1}{2\theta} \int_{\mathbb{R}^+ \times \mathbb{R}^d} \frac{d\lambda}{\lambda^{1-\alpha_0}} \mu(d\xi) \frac{\theta + \lambda}{(\theta + \lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2} \leq C \theta^{-2}.$$

Notice that

$$\frac{1}{(\theta + \lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2} \leq \frac{1}{\theta} \frac{\theta + \lambda}{(\theta + \lambda)^2 + |\xi|^2} \frac{1}{\theta^2 + |\xi|^2}.$$

The first term in (A.2) yields the same bound. $\square$

Funding. The author was supported in part by the Simons Foundation #585506.

REFERENCES

- [1] Anderson, P. W. Localized Magnetic States in Metals. *Phys. Rev.* **124** (1961) 41–53.
- [2] Balan, R. M. Stratonovich solution for the wave equation. *J. Theoret. Probab.* **35** (2022), 2643–2689.
- [3] Balan, R. M. and Conus, D. Intermittency for wave and heat equations with fractional noise in time. *Ann. Probab.* **44** (2016) 1488–1534.
- [4] Balan, R. M., Chen, L. and Chen, X. Exact asymptotics of the stochastic wave equation with time-independent noise. *Annales de l’Institut Henri Poincaré* **58** (2022), 1590–1620.
- [5] Chen, X. *Random Walk Intersections: Large Deviations and Related Topics*. Mathematical Surveys and Monographs, **157**. American Mathematical Society, Providence 2009.
- [6] Chen, X. Quenched asymptotics for Brownian motion in generalized Gaussian potential. *Ann. Probab.* **42** (2014), 576–622.
- [7] Chen, X. Spatial asymptotics for the parabolic Anderson models with generalized time-space Gaussian noise. *Ann. Probab.* **44** (2016), 1535–1598.
- [8] Chen, X. Moment asymptotics for parabolic Anderson equation with fractional time-space noise: in Skorokhod regime *Annales de l’Institut Henri Poincaré* **53** (2017) 819–841.
- [9] Chen, X. Parabolic Anderson model with rough or critical Gaussian noise. *Annales de l’Institut Henri Poincaré* **55** (2019), 941–976.
- [10] Chen, X. Exponential asymptotics for Brownian self-intersection local times under Dalang’s condition. *Electronic J. P.* **28** (2023), 1–17.
- [11] Chen, X., Deya, A. Song, J. and Tindel, S. Hyperbolic Anderson model 2: Strichartz estimates and Stratonovich setting *PDF International Mathematics Research Notices*. **00** (2023), 1–54.
- [12] Chen, X., Deya, A. Song, J. and Tindel, S. Solving the hyperbolic Anderson model I: Skorokhod setting. *Annales de l’Institut Henri Poincaré* **61** (2025), 1794–1814.
- [13] Chen, X., Hu, Y. Song, J. and Xing, F. Exponential asymptotics for time-space Hamiltonians. *Annales de l’Institut Henri Poincaré* **51** (2015), 1529–1561.
- [14] Chen, X. and Hu, Y. Hyperbolic Anderson equations with general time-independent Gaussian noise: Stratonovich regime *Ann. Probab.* **53** (2025) 2144–2195.
- [15] Dalang, R. C. Extending martingale measure stochastic integral with applications to spatially homogeneous S.P.D.E’s. *Electron. J. Probab.* **4** (1999), 1–29.
- [16] Dalang, R. C., Mueller, C. and Tribe, R. A Feynman-Kac-type formula for the deterministic and stochastic wave equations and other P.D.E.’s. *Trans. Amer. Math. Soc.* **360** (2008), 4681–4703.
- [17] Dembo, A. and Zeitouni, O. (1997). *Large Deviations Techniques and Applications*. 2nd ed. Springer, New York.
- [18] Gärtner, J. and Molchanov, S. Moment asymptotics and Lifshitz tails for the parabolic Anderson model. *Stochastic models (Ottawa, ON, 1998)*, 141–157, *CMS Conf. Proc.*, 26, Amer. Math. Soc., Providence, RI, 2000.

- [19] Hu, Y. *Analysis on Gaussian spaces*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2017.
- [20] Hu, Y. Z. and Meyer, P.-A. Sur les intégrales multiples de Stratonovitch. *Séminaire de Probabilités, XXII*, 72–81, *Lecture Notes in Math.*, 1321, Springer, Berlin, 1988.
- [21] Hu, Y. and Meyer, P. A. On the approximation of multiple Stratonovich integrals. *Stochastic processes*, 141-147, Springer, New York, 1993.
- [22] Hu, Y., Huang, J., Nualart, D. and Tindel, S. Stochastic heat equations with general multiplicative Gaussian noise: Hölder continuity and intermittency. *Electron. J. Probab.* **20** (2015) 1-50
- [23] Evans, L. C. *Partial differential equations*. Second edition. *Graduate Studies in Mathematics*, 19. American Mathematical Society, Providence, RI, 2010.
- [24] Feller, W. (1971). *An introduction to Probability Theory and Its Applications II* 2nd ed. Wiley, New York.
- [25] Hairer, M. Solving the KPZ equation. *Ann. of Math.* **178** (2013) 559–664.
- [26] König, W. *The parabolic Anderson Model. Random Walk in random potential. Pathways in Mathematics*. Birkhäuser/Springer, [Cham], 2016.
- [27] Kallenberg, O. (2002). *Foundations of Modern Probability*, 2nd ed. Springer, New York.
- [28] Kardar, M., Parisi, G. and Zhang, Y.-C. Dynamic Scaling of Growing Interfaces. *Physical Review Letters*. **56** (9), 1986, 889–892.
- [29] Li, W.V. and Shao, Q.-M. Gaussian processes: inequalities, small ball probabilities and applications. *Handbook Statist.* **19** (2001), 533–597.
- [30] Lipschutz, S., Spiegel, M. R. and Liu, J. *Mathematical Handbook of Formulas and Tables*, Schaum's Outline Series (3rd ed.), (2009) McGraw-Hill.
- [31] Marcus, M. B. and Rosen, J. *Markov Processes, Gaussian Processes, and Local Times* *Cambridge Studies in Advanced Mathematics* **100**, Cambridge Univ. Press, Cambridge (2006)
- [32] Pipiras, V. and Taqqu, M. S. Integration questions related to fractional Brownian motion. *Probab. Theory Related Fields* **118** (2000), no. 2, 251-291.
- [33] Song, J. On a class of stochastic partial differential equations. *Stochastic Process. Appl.* **127** (2017), 37–79.